

Moments, intermittency and growth indices for the nonlinear fractional stochastic heat equation

Le Chen & Robert C. Dalang

**Stochastic Partial Differential
Equations: Analysis and
Computations**

ISSN 2194-0401
Volume 3
Number 3

Stoch PDE: Anal Comp (2015) 3:360-397
DOI 10.1007/s40072-015-0054-x

Volume 3 • Number 3 September 2015

Stochastic Partial Differential Equations Analysis and Computations

Editors in Chief
A. Debussche
École Normale Supérieure de Rennes, France
B. Rozovsky
Brown University, USA

Editorial Board

D. Crisan <i>Imperial College London, UK</i>	S. Kuksin <i>École Polytechnique, Paris, France</i>	G. Papanicolaou <i>Stanford University, USA</i>
R. Dalang <i>École Polytechnique Fédérale de Lausanne, Switzerland</i>	S. Lototsky <i>University of Southern California, USA</i>	E. Pardoux <i>University of Marseille, France</i>
G. Da Prato <i>Scuola Normale Superiore, Pisa, Italy</i>	T. Lyons <i>University of Oxford, UK</i>	M. Roekner <i>University of Bielefeld, Germany</i>
A. De Bouard <i>École Polytechnique, Paris, France</i>	J. Mattingly <i>Duke University, USA</i>	M. Sanz-Solé <i>University of Barcelona, Spain</i>
F. Flandoli <i>University of Pisa, Italy</i>	S. Méléard <i>École Polytechnique, Paris, France</i>	C. Schwab <i>ETH Zurich, Switzerland</i>
T. Funaki <i>University of Tokyo, Japan</i>	R. Mikulevicius <i>University of Southern California, USA</i>	P. Souganidis <i>University of Chicago, USA</i>
I. Gyöngy <i>University of Edinburgh, UK</i>	C. Mueller <i>University of Rochester, USA</i>	A. Stuart <i>University of Warwick, UK</i>
M. Hairer <i>University of Warwick, UK</i>	F. Nobile <i>École Polytechnique Fédérale de Lausanne, Switzerland</i>	
G. Karniadakis <i>Brown University, USA</i>	D. Nualart <i>University of Kansas, USA</i>	

 Springer
40072 • 3(3) 291–444 (2015)
ISSN 2194-0401 (print)
ISSN 2194-041X (electronic)

Your article is protected by copyright and all rights are held exclusively by Springer Science +Business Media New York. This e-offprint is for personal use only and shall not be self-archived in electronic repositories. If you wish to self-archive your article, please use the accepted manuscript version for posting on your own website. You may further deposit the accepted manuscript version in any repository, provided it is only made publicly available 12 months after official publication or later and provided acknowledgement is given to the original source of publication and a link is inserted to the published article on Springer's website. The link must be accompanied by the following text: "The final publication is available at link.springer.com".

Moments, intermittency and growth indices for the nonlinear fractional stochastic heat equation

Le Chen^{1,2} · Robert C. Dalang²

Received: 12 September 2014 / Published online: 10 July 2015
© Springer Science+Business Media New York 2015

Abstract We study the nonlinear fractional stochastic heat equation in the spatial domain \mathbb{R} driven by space-time white noise. The initial condition is taken to be a measure on \mathbb{R} , such as the Dirac delta function, but this measure may also have non-compact support. Existence and uniqueness, as well as upper and lower bounds on all p th moments ($p \geq 2$), are obtained. These bounds are uniform in the spatial variable, which answers an open problem mentioned in Conus and Khoshnevisan (Probab Theory Relat Fields 152(3–4):681–701, 2012). We improve the weak intermittency statement by Foondun and Khoshnevisan (Electron J Probab 14(21):548–568, 2009), and we show that the growth indices (of linear type) introduced in Conus and Khoshnevisan (Probab Theory Relat Fields 152(3–4):681–701, 2012) are infinite. We introduce the notion of “growth indices of exponential type” in order to characterize the manner in which high peaks propagate away from the origin, and we show that the presence of a fractional differential operator leads to significantly different behavior compared with the standard stochastic heat equation.

Keywords Nonlinear fractional stochastic heat equation · Parabolic Anderson model · Rough initial data · Intermittency · Growth indices · Stable processes

Le Chen and Robert C. Dalang were supported in part by the Swiss National Foundation for Scientific Research.

✉ Le Chen
chenle02@gmail.com

Robert C. Dalang
robert.dalang@epfl.ch

¹ Present Address: Department of Mathematics, University of Kansas, 405 Snow Hall, 1460 Jayhawk Blvd, Lawrence, KS 66045-7594, USA

² Institut de mathématiques, École Polytechnique Fédérale de Lausanne, Station 8, 1015 Lausanne, Switzerland

Mathematics Subject Classification Primary 60H15 · Secondary 60G60 · 35R60

1 Introduction

In this paper, we consider the following nonlinear fractional stochastic heat equation:

$$\begin{cases} \left(\frac{\partial}{\partial t} - {}_x D_\delta^a \right) u(t, x) = \rho(u(t, x)) \dot{W}(t, x), & t \in \mathbb{R}_+^* :=]0, +\infty[, x \in \mathbb{R}, \\ u(0, \circ) = \mu(\circ), \end{cases} \quad (1.1)$$

where $a \in]0, 2]$ is the order of the fractional differential operator ${}_x D_\delta^a$ and δ ($|\delta| \leq a \wedge (2 - a) := \min(a, 2 - a)$) is its skewness, \dot{W} is the space-time white noise, μ is the initial data (a measure), the function $\rho : \mathbb{R} \mapsto \mathbb{R}$ is Lipschitz continuous, and \circ denotes the spatial dummy variable. The definition and properties of the fractional differential operator ${}_x D_\delta^a$ are recalled in Sect. 2.1. This equation falls into a class of equations studied by Debbi and Dozzi [12, 13]. According to [11, Theorem 11], even the linear form of (1.1) ($\rho \equiv 1$) does not have a solution if $a \leq 1$, so they consider $a \in]1, 2]$. If we focus on deterministic initial conditions, then in the setting of (1.1), they proved in [13, Theorem 1] that there is a unique random field solution if μ has a bounded density.

Equation (1.1) is of particular interest since it is an extension of the classical *parabolic Anderson model* [5], in which $a = 2$ and $\delta = 0$, so ${}_x D_\delta^a$ is the operator $\partial^2 / \partial x^2$, and $\rho(u) = \lambda u$ is a linear function. Foondun and Khoshnevisan [15] considered problem (1.1) with the operator ${}_x D_\delta^a$ replaced by the $L^2(\mathbb{R})$ -generator \mathcal{L} of a Lévy process. They proved the existence of a random field solution under the assumption that the initial data μ has a bounded and nonnegative density. In [9], the operator ${}_x D_\delta^a$ is replaced by the generator of a symmetric Lévy process and the authors prove that μ can be any finite Borel measure on \mathbb{R} . Recently, Balan and Conus [1, 2] studied the Anderson model with fractional Laplacian and bounded initial condition, and with Gaussian, spatially homogeneous noise that behaves in time like a fractional Brownian motion with Hurst index $H > 1/2$.

Following the approach of [6], in which the case $a = 2$ and $\delta = 0$ was considered, we begin by extending the above results (for the operator ${}_x D_\delta^a$) to allow a wider class of initial data: Let $\mathcal{M}(\mathbb{R})$ be the set of signed Borel measures on \mathbb{R} . From the Jordan decomposition, $\mu = \mu_+ - \mu_-$ where μ_\pm are two non-negative Borel measures with disjoint support, and denote $|\mu| = \mu_+ + \mu_-$. Then our admissible initial data is $\mu \in \mathcal{M}_a(\mathbb{R})$, where

$$\mathcal{M}_a(\mathbb{R}) := \left\{ \mu \in \mathcal{M}(\mathbb{R}) : \sup_{y \in \mathbb{R}} \int_{\mathbb{R}} |\mu|(dx) \frac{1}{1 + |x - y|^{1+a}} < +\infty \right\}, \quad \text{for } a \in]1, 2].$$

We will also use the set $\mathcal{M}_{a,+}(\mathbb{R}) = \{ \mu \in \mathcal{M}_a(\mathbb{R}) : \mu \text{ is non negative} \}$. For $\mu \in \mathcal{M}_a(\mathbb{R})$, we obtain estimates for the moments $\mathbb{E}(|u(t, x)|^p)$ for all $p \geq 2$. These estimates have the same structure as those that are given in [6, Theorem 2.4], but the kernel \mathcal{K} that appears in this reference has quite different properties than those of the kernel \mathcal{K} that appears in relation with Eq. (1.1): see Sect. 3.2.

Let us define the *upper and lower Lyapunov exponents of order p* by

$$\bar{m}_p(x) := \limsup_{t \rightarrow +\infty} \frac{1}{t} \log \mathbb{E} (|u(t, x)|^p), \quad \underline{m}_p(x) := \liminf_{t \rightarrow +\infty} \frac{1}{t} \log \mathbb{E} (|u(t, x)|^p), \tag{1.2}$$

for all $p \geq 2$ and $x \in \mathbb{R}$. If the initial data is constant, then \underline{m}_p and \bar{m}_p do not depend on x . In this case, the solution is called *fully intermittent* if $\underline{m}_2 > 0$ and $m_1 = 0$ by Carmona and Molchanov [5, Definition III.1.1, on p. 55]. For a detailed discussion of the meaning of this intermittency property, see [17]. Informally, it means that the sample paths of $u(t, x)$ exhibit “high peaks” separated by “large valleys”.

Foondun and Khoshnevisan proved *weak intermittency* in [15], namely, for all $p \geq 2$,

$$\bar{m}_2(x) > 0, \quad \text{and} \quad \bar{m}_p(x) < +\infty \quad \text{for all } x \in \mathbb{R},$$

under the conditions that $\mu(dx) = f(x)dx$ with $\inf_{x \in \mathbb{R}} f(x) > 0$ and $\inf_{x \neq 0} |\rho(x)/x| > 0$. We improve this result by showing in Theorem 3.4 that when $1 < a < 2$, $|\delta| < 2 - a$ (strict inequality) and $\mu \in \mathcal{M}_a(\mathbb{R})$ is nonnegative and nonvanishing, then for all $p \geq 2$,

$$\inf_{x \in \mathbb{R}} \underline{m}_p(x) > 0, \quad \text{and} \quad \sup_{x \in \mathbb{R}} \bar{m}_p(x) < +\infty.$$

For this, we need a growth condition on ρ , namely, that for some constants $l_\rho > 0$ and $\underline{\zeta} \geq 0$,

$$\rho(x)^2 \geq l_\rho^2 (\underline{\zeta}^2 + x^2), \quad \text{for all } x \in \mathbb{R}. \tag{1.3}$$

In a forthcoming paper [8], this weak intermittency property will be extended to full intermittency by showing in addition that $m_1(x) \equiv 0$.

Our result answers an open problem stated by Conus and Khoshnevisan [10]. Indeed, for the case of the fractional Laplacian, which corresponds to our setting with $a \in]1, 2[$ and $\delta = 0$, they ask whether the function $t \mapsto \sup_{x \in \mathbb{R}} \mathbb{E} (|u(t, x)|^2)$ has exponential growth in t for initial data with exponential decay. Our answer is “yes” under the condition (1.3). In addition, under these conditions, if $\mu \in \mathcal{M}_{a,+}(\mathbb{R})$ and $\mu \neq 0$, then for fixed $x \in \mathbb{R}$, the function $t \mapsto \mathbb{E} (|u(t, x)|^2)$ has at least exponential growth; see Remark 3.5.

We define the following *growth indices of exponential type*:

$$\underline{e}(p) := \sup \left\{ \alpha > 0 : \lim_{t \rightarrow \infty} \frac{1}{t} \sup_{|x| \geq \exp(\alpha t)} \log \mathbb{E} (|u(t, x)|^p) > 0 \right\}, \tag{1.4}$$

$$\bar{e}(p) := \inf \left\{ \alpha > 0 : \lim_{t \rightarrow \infty} \frac{1}{t} \sup_{|x| \geq \exp(\alpha t)} \log \mathbb{E} (|u(t, x)|^p) < 0 \right\}, \tag{1.5}$$

in order to give a proper characterization of the propagation speed of “high peaks”. This concept is discussed in Conus and Khoshnevisan [10]. These authors define analogous

indices $\underline{\lambda}(p)$ and $\bar{\lambda}(p)$, in which $|x| \geq \exp(\alpha t)$ is replaced by $|x| \geq \alpha t$, which we call *growth indices of linear type*.

Conus and Khoshnevisan [10] consider the case where ${}_x D_\delta^a$ is replaced by the generator \mathcal{L} of a real-valued symmetric Lévy process $(X_t, t \geq 0)$. They showed in [10, Theorem 1.1 and Remark 1.2] that if the initial data μ is a nonnegative lower semicontinuous function with certain exponential decay at infinity, and if X_1 has exponential moments, then

$$0 < \underline{\lambda}(p) \leq \bar{\lambda}(p) < +\infty, \quad \text{for all } p \in [2, +\infty) .$$

For example, a Lévy process that satisfies this assumption is the “truncated symmetric stable process”.

Among Lévy processes, stable processes constitute an important subclass with a self-similarity property. Infinitesimal generators of these processes are not covered by the results of [10], since even the second moment of X_1 does not exist. It turns out that the fractional differential operator ${}_x D_\delta^a$ is the infinitesimal generator of a (not necessarily symmetric) strictly stable process with $a \in]1, 2]$ (see Sect. 2.1), and we will see that when $1 < a < 2$, the presence of the fractional differential operator ${}_x D_\delta^a$ in (1.1) leads to significantly different behaviors of the speed of propagation of high peaks, compared to that obtained in [10].

Indeed, we show first that if the initial data has sufficiently rapid decay at $\pm\infty$, then $\bar{e}(p) < \infty$ (see (3.22) and (3.23)). Then we show that if $1 < a < 2$ and $|\delta| < 2 - a$ (meaning that the underlying stable process has both positive and negative jumps), then

$$\underline{e}(p) > 0, \quad \text{for all } p \in [2, +\infty) \text{ and } \mu \in \mathcal{M}_{a,+}(\mathbb{R}), \mu \neq 0, \quad (1.6)$$

provided ρ satisfies condition (1.3). This conclusion applies, for instance, to the case where the initial data μ is the Dirac delta function. In particular, for well-localized initial data (for instance, $\mu \geq 0$ and $\int_{\mathbb{R}} |y|^\eta \mu(dy) < \infty$ for some $\eta > 0$), $0 < \underline{e}(p) \leq \bar{e}(p) < +\infty$, whereas for initial data that is bounded below ($\mu(dx) = f(x)dx$ with $f(x) > c > 0$, for all $x \in \mathbb{R}$), $\underline{e}(p) = \bar{e}(p) = +\infty$. See Theorem 3.6 for the precise statements. As a direct consequence, $\underline{\lambda}(p) = \bar{\lambda}(p) = +\infty$ for all $p \in [2, \infty[$.

The structure of this paper is as follows. After defining the operator ${}_x D_\delta^a$ and giving the meaning of (1.1) in Sect. 2, the main results are presented in Sect. 3: Existence and general bounds are given in Theorem 3.1. These bounds are expressed in terms of the kernel \mathcal{K} mentioned above, for which explicit upper and lower bounds are given. These lead to our results on weak intermittency (Theorem 3.4) and growth indices (Theorem 3.6). Section 4 contains the proof of Theorem 3.1 and Sect. 5 presents the proofs of Theorems 3.4 and 3.6.

2 Preliminaries and notation

We begin by defining the differential operator ${}_x D_\delta^a$ that appears in the spde (1.1), then we shall give the rigorous meaning of the spde.

2.1 The Riesz–Feller fractional derivative

Let $\mathcal{F}f(\xi) = \int_{\mathbb{R}} dx e^{-i\xi x} f(x)$ denote the Fourier transform. For $0 < a \leq 2$ and $|\delta| \leq \min(a, 2 - a)$, the Riesz–Feller fractional derivative ${}_x D_\delta^a f$ of a smooth and integrable function f is defined (see [19, (2.2)]) by

$$\mathcal{F}({}_x D_\delta^a f)(\xi) = {}_\delta \psi_a(\xi) \mathcal{F}f(\xi), \quad \text{where } {}_\delta \psi_a(\xi) = -|\xi|^a \exp(-i\pi \delta \operatorname{sgn}(\xi)/2). \tag{2.1}$$

When $a = 2$ (and therefore $\delta = 0$), this is simply the ordinary second derivative $\frac{d^2}{dx^2}$. For $1 < a < 2$ and $\delta \leq 2 - a$, which is the case that we are most interested in, as stated in [19, (2.8)] (in which integrals are understood in the sense of Cauchy principle values), this is equivalent to the more explicit formula

$$\begin{aligned} {}_x D_\delta^a f(x) &= c_a^+ \int_0^{+\infty} \frac{f(x+z) - f(x) - zf'(x)}{z^{1+a}} dz \\ &\quad + c_a^- \int_{-\infty}^0 \frac{f(x+z) - f(x) - zf'(x)}{(-z)^{1+a}} dz, \end{aligned} \tag{2.2}$$

where

$$c_a^\pm = \frac{\Gamma(1+a)}{\pi} \sin\left((a \pm \delta)\frac{\pi}{2}\right)$$

and $\Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt$ is Euler’s Gamma function [21]. Indeed, taking the Fourier transform of the right-hand side of (2.2) leads to (2.1) by (2.3) below and elementary properties of Fourier transform.

From the point of view of a probabilist, the operator ${}_x D_\delta^a$ is the infinitesimal generator of a strictly a -stable Lévy process $X = (X_t, t \geq 0)$ (where “strictly” refers to the fact that the process is centered: see [23, Chapter 3]), with Lévy measure

$$\nu_a(dz) = c_a^+ \frac{dz}{z^{1+a}} 1_{\{z>0\}} + c_a^- \frac{dz}{(-z)^{1+a}} 1_{\{z<0\}}.$$

Indeed, the general form of the infinitesimal generator of X , given in terms of its Lévy measure in [23, Theorem 31.5, p. 208], is

$$\begin{aligned} \mathcal{L}f(x) &= c_a^+ \int_0^{+\infty} \frac{f(x+z) - f(x) - zf'(x)}{z^{1+a}} dz \\ &\quad + c_a^- \int_{-\infty}^0 \frac{f(x+z) - f(x) - zf'(x)}{(-z)^{1+a}} dz, \end{aligned}$$

where no “truncation function” nor additional drift term appears because of the centering, and the characteristic function of X_t , given by the Lévy–Khintchine formula [23, Theorem 8.1], is

$$\exp \left[-t \int_{-\infty}^{+\infty} (e^{i\xi z} - 1 - i\xi z) \nu_a(dz) \right] = \exp(-t \delta \psi_a(\xi)), \tag{2.3}$$

where the right-hand side can be obtained from the left-hand side via a direct calculation using the formula $\Gamma(a)\Gamma(1 - a) = \pi / \sin(\pi a)$ [21, 5.5.3, p. 138] and the identity

$$\int_0^{+\infty} \frac{e^{-qz} - 1 + qz}{z^{1+a}} dz = q^a \Gamma(-a), \quad q \in \mathbb{C}, \quad 1 < \Re(a) < 2,$$

which follows from [21, 5.9.5, p. 140].

In order to study the spde (1.1), we need the fundamental solution of the operator $\frac{\partial}{\partial t} - {}_x D_\delta^a$. According to the above discussion, this is

$$\delta G_a(t, x) := \mathcal{F}^{-1} \left[\exp \{ \delta \psi_a(\cdot) t \} \right] (x) = \frac{1}{2\pi} \int_{\mathbb{R}} d\xi \exp \left\{ i\xi x - t |\xi|^a e^{-i\pi \delta \operatorname{sgn}(\xi)/2} \right\}, \tag{2.4}$$

where \mathcal{F}^{-1} is the inverse Fourier transform. This is (one way to represent) the density of the strictly a -stable random variable X_t with Lévy measure $t\nu_a$, as given in the representation (C) in [27, p. 17]. For comparison, the expression in [23, Theorem 14.15] corresponds to the representation (A) in [27, p. 9]. For strictly stable laws, in particular the case $1 < a < 2$, these two representations are equivalent: see [27, (I.28),(I.26) and Theorem C.3]. Properties of δG_a will be given in Lemma 4.1 below. We refer to [19] for more details on these fractional differential operators.

In this paper, we denote the solution to the homogeneous equation

$$\begin{cases} \left(\frac{\partial}{\partial t} - {}_x D_\delta^a \right) u(t, x) = 0, & t \in \mathbb{R}_+^*, \quad x \in \mathbb{R}, \\ u(0, \circ) = \mu(\circ), \end{cases}$$

by

$$J_0(t, x) := (\delta G_a(t, \circ) * \mu) (x) = \int_{\mathbb{R}} \mu(dy) \delta G_a(t, x - y),$$

where “ $*$ ” denotes the convolution in the space variable.

2.2 The stochastic pde

Let $W = \{W_t(A), A \in \mathcal{B}_b(\mathbb{R}), t \geq 0\}$ be a space-time white noise defined on a probability space (Ω, \mathcal{F}, P) , where $\mathcal{B}_b(\mathbb{R})$ is the collection of Borel sets with finite Lebesgue measure. Let $(\mathcal{F}_t, t \geq 0)$ be the filtration generated by W and augmented by the σ -field \mathcal{N} generated by all P -null sets in \mathcal{F} :

$$\mathcal{F}_t = \sigma (W_s(A) : 0 \leq s \leq t, A \in \mathcal{B}_b(\mathbb{R})) \vee \mathcal{N}, \quad t \geq 0.$$

In the following, we fix this filtered probability space $\{\Omega, \mathcal{F}, (\mathcal{F}_t, t \geq 0), P\}$. We use $\|\cdot\|_p$ to denote the $L^p(\Omega)$ -norm ($p \geq 1$). With this setup, W becomes a worthy martingale measure in the sense of Walsh [25], and $\iint_{[0,t] \times \mathbb{R}} X(s, y)W(ds, dy)$ is well-defined in this reference for a suitable class of random fields $\{X(s, y), (s, y) \in \mathbb{R}_+ \times \mathbb{R}\}$.

The rigorous meaning of the spde (1.1) uses the integral formulation

$$u(t, x) = J_0(t, x) + I(t, x), \quad \text{where} \tag{2.5}$$

$$I(t, x) = \iint_{[0,t] \times \mathbb{R}} \delta G_a(t - s, x - y) \rho(u(s, y)) W(ds, dy).$$

Definition 2.1 A process $u = (u(t, x), (t, x) \in \mathbb{R}_+^* \times \mathbb{R})$ is called a *random field solution* to (1.1) if:

- (1) u is adapted, i.e., for all $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$, $u(t, x)$ is \mathcal{F}_t -measurable;
- (2) u is jointly measurable with respect to $\mathcal{B}(\mathbb{R}_+^* \times \mathbb{R}) \times \mathcal{F}$;
- (3) $(\delta G_a^2 \star \|\rho(u)\|_2^2)(t, x) < +\infty$ for all $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$, where “ \star ” denotes the simultaneous convolution in both space and time variables, that is,

$$\left(\delta G_a^2 \star \|\rho(u)\|_2^2\right)(t, x) := \int_0^t ds \int_{\mathbb{R}} dy \delta G_a^2(t - s, x - y) \|\rho(u(s, y))\|_2^2;$$

- (4) For all $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$, $u(t, x)$ satisfies (2.5) a.s.;
- (5) The function $(t, x) \mapsto I(t, x)$ mapping $\mathbb{R}_+^* \times \mathbb{R}$ into $L^2(\Omega)$ is continuous.

Assume that the function $\rho : \mathbb{R} \mapsto \mathbb{R}$ is globally Lipschitz continuous with Lipschitz constant $\text{Lip}_\rho > 0$. This implies a growth condition on ρ : for some constants $L_\rho > 0$ and $\bar{\varsigma} \geq 0$,

$$\rho(x)^2 \leq L_\rho^2 \left(\bar{\varsigma}^2 + x^2\right), \quad \text{for all } x \in \mathbb{R}. \tag{2.6}$$

Note that one can take $\bar{\varsigma} = |\rho(0)|$ and $L_\rho \leq \sqrt{2} \text{Lip}_\rho$ (and the inequality may even be strict). We shall also specially consider the linear case $\rho(u) = \lambda u$ with $\lambda \neq 0$, which is related to the *parabolic Anderson model* ($a = 2$). It is a special case of the following near-linear growth condition: for some constant $\varsigma \geq 0$,

$$\rho(x)^2 = \lambda^2 \left(\varsigma^2 + x^2\right), \quad \text{for all } x \in \mathbb{R}. \tag{2.7}$$

For all $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$, $n \in \mathbb{N}$ and $\lambda \in \mathbb{R}$, define

$$\mathcal{L}_0(t, x; \lambda) := \lambda^2 \delta G_a^2(t, x),$$

$$\mathcal{L}_n(t, x; \lambda) := \underbrace{(\mathcal{L}_0 \star \dots \star \mathcal{L}_0)}_{n+1 \text{ factors } \mathcal{L}_0(\cdot, \cdot; \lambda)}(t, x; \lambda), \quad \text{for } n \geq 1, \tag{2.8}$$

and

$$\mathcal{K}(t, x; \lambda) := \sum_{n=0}^{\infty} \mathcal{L}_n(t, x; \lambda) \tag{2.9}$$

(the convergence of this series is established in Proposition 3.2). For $t \geq 0$, define

$$\mathcal{H}(t; \lambda) := (1 \star \mathcal{K}(\cdot, \circ; \lambda))(t, x)$$

(notice that the right-hand side does not depend on x).

Let z_p be the universal constant in the Burkholder-Davis-Gundy inequality (in particular, $z_2 = 1$), and so $z_p \leq 2\sqrt{p}$ for all $p \geq 2$; see [4, Appendix]. Define

$$a_{p, \bar{\zeta}} = \begin{cases} 2^{(p-1)/p} & \text{if } \bar{\zeta} \neq 0 \text{ and } p > 2, \\ \sqrt{2} & \text{if } \bar{\zeta} = 0 \text{ and } p > 2, \\ 1 & \text{if } p = 2. \end{cases}$$

We apply the following conventions to the kernel functions $\mathcal{K}(t, x; \lambda)$ (and similarly to $\mathcal{H}(t; \lambda)$):

$$\begin{aligned} \mathcal{K}(t, x) &:= \mathcal{K}(t, x; \lambda), & \bar{\mathcal{K}}(t, x) &:= \mathcal{K}(t, x; L_\rho), \\ \underline{\mathcal{K}}(t, x) &:= \mathcal{K}(t, x; l_\rho), & \widehat{\mathcal{K}}_p(t, x) &:= \mathcal{K}(t, x; a_{p, \bar{\zeta}} z_p L_\rho), \end{aligned} \quad \text{for } p \geq 2.$$

3 Main results

3.1 Existence, uniqueness and moments

The following theorem extends the result of [6, Theorem 2.4] from $a = 2$ to $a \in]1, 2]$. In view of the related result [7, Theorem 2.3] and Remark 2.4 in this reference, the bounds in this theorem are not a surprise, though they do require a proof. The main effort will be to turn these abstract bounds into concrete estimates, via explicit upper and lower bounds on the functions \mathcal{K} and \mathcal{H} (see Sect. 3.2). For $\tau \geq t > 0$ and $x, y \in \mathbb{R}$, define

$$\begin{aligned} \mathcal{I}(t, x, \tau, y; \zeta, \lambda) &:= \lambda^2 \int_0^t dr \int_{\mathbb{R}} dz \left[J_0^2(r, z) + \left(J_0^2(\cdot, \circ) \star \mathcal{K}(\cdot, \circ; \lambda) \right)(r, z) \right. \\ &\quad \left. + \zeta^2 (\mathcal{H}(r; \lambda) + 1) \right] \delta G_a(t - r, x - z) \delta G_a(\tau - r, y - z). \end{aligned}$$

Theorem 3.1 (Existence, uniqueness and moments) *Suppose that*

- (i) $1 < a \leq 2$ and $|\delta| \leq 2 - a$;
- (ii) the function ρ is Lipschitz continuous and satisfies the growth condition (2.6);
- (iii) the initial data are such that $\mu \in \mathcal{M}_a(\mathbb{R})$.

Then the stochastic pde (1.1) has a random field solution $(u(t, x), (t, x) \in \mathbb{R}_+^* \times \mathbb{R})$. Moreover:

- (1) $u(t, x)$ is unique in the sense of versions;
- (2) $(t, x) \mapsto u(t, x)$ is $L^p(\Omega)$ -continuous for all integers $p \geq 2$;
- (3) For all even integers $p \geq 2$, all $\tau \geq t > 0$ and $x, y \in \mathbb{R}$,

$$\|u(t, x)\|_p^2 \leq \begin{cases} J_0^2(t, x) + ([\bar{\zeta}^2 + J_0^2] \star \bar{\mathcal{K}})(t, x), & \text{if } p = 2, \\ 2J_0^2(t, x) + ([\bar{\zeta}^2 + 2J_0^2] \star \widehat{\mathcal{K}}_p)(t, x), & \text{if } p > 2, \end{cases} \quad (3.1)$$

and

$$\mathbb{E}[u(t, x)u(\tau, y)] \leq J_0(t, x)J_0(\tau, y) + \mathcal{I}(t, x, \tau, y; \bar{\zeta}, L_\rho). \quad (3.2)$$

- (4) If ρ satisfies (1.3), then for all $\tau \geq t > 0$ and $x, y \in \mathbb{R}$,

$$\|u(t, x)\|_2^2 \geq J_0^2(t, x) + \left((\underline{\zeta}^2 + J_0^2) \star \underline{\mathcal{K}} \right)(t, x), \quad (3.3)$$

and

$$\mathbb{E}[u(t, x)u(\tau, y)] \geq J_0(t, x)J_0(\tau, y) + \mathcal{I}(t, x, \tau, y; \underline{\zeta}, l_\rho). \quad (3.4)$$

- (5) If ρ satisfies (2.7), then for all $\tau \geq t > 0$ and $x, y \in \mathbb{R}$,

$$\|u(t, x)\|_2^2 = J_0^2(t, x) + \left((\zeta^2 + J_0^2) \star \mathcal{K} \right)(t, x), \quad (3.5)$$

and

$$\mathbb{E}[u(t, x)u(\tau, y)] = J_0(t, x)J_0(\tau, y) + \mathcal{I}(t, x, \tau, y; \zeta, \lambda). \quad (3.6)$$

The proof of this theorem is explained in Sect. 4.

3.2 Estimates on the kernel function $\mathcal{K}(t, x)$

Recall that if the partial differential operator is the heat operator $\frac{\partial}{\partial t} - \frac{\nu}{2} \Delta$, then

$$\mathcal{K}^{\text{heat}}(t, x; \lambda) = G_{\frac{\nu}{2}}(t, x) \left(\frac{\lambda^2}{\sqrt{4\pi\nu t}} + \frac{\lambda^4}{2\nu} e^{\frac{\lambda^4 t}{4\nu}} \Phi \left(\lambda^2 \sqrt{\frac{t}{2\nu}} \right) \right), \quad (3.7)$$

where $\nu > 0$ and $\Phi(x)$ is the distribution function of a standard Normal random variable; see [6, Proposition 2.2]. When the partial differential operator is the wave operator $\frac{\partial^2}{\partial t^2} - \kappa^2 \Delta$,

$$\mathcal{K}^{\text{wave}}(t, x; \lambda) = \frac{\lambda^2}{4} I_0 \left(\sqrt{\frac{\lambda^2((\kappa t)^2 - x^2)}{2\kappa}} \right) 1_{\{|x| \leq \kappa t\}}, \tag{3.8}$$

where $\kappa > 0$ and $I_0(x)$ is the modified Bessel function of the first kind of order 0; see [7, Proposition 3.1].

Except in the above two cases, we do not have an explicit formula for the kernel function $\mathcal{K}(t, x)$ in (2.9). In order to make use of the moment formulas in (3.1) and (3.3), we derive upper and lower bounds on this kernel function in the following two propositions. We will need the two-parameter *Mittag-Leffler function* [22, Sect. 1.2]:

$$E_{\alpha, \beta}(z) := \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}, \quad \alpha > 0, \beta > 0. \tag{3.9}$$

Let a^* be the dual of a : $1/a + 1/a^* = 1$. By Lemma 4.1 below (Property (ii)), the constant

$$\Lambda = {}_{\delta} \Lambda_a := \sup_{x \in \mathbb{R}} {}_{\delta} G_a(1, x) \tag{3.10}$$

is finite. In particular,

$${}_0 \Lambda_a = {}_0 G_a(1, 0) = \frac{1}{2\pi} \int_{\mathbb{R}} d\xi \exp(-|\xi|^a) = \frac{1}{a\pi} \int_0^{\infty} dt e^{-t} t^{-1+1/a} = \frac{\Gamma(1 + 1/a)}{\pi}.$$

In the following, we often omit the dependence of ${}_{\delta} \Lambda_a$ on δ and a and simply write Λ instead of ${}_{\delta} \Lambda_a$. Define

$$\begin{aligned} \gamma &:= \lambda^2 \Lambda \Gamma(1/a^*), & \bar{\gamma} &:= L_{\rho}^2 \Lambda \Gamma(1/a^*), \\ \underline{\gamma} &:= l_{\rho}^2 \Lambda \Gamma(1/a^*), & \hat{\gamma}_p &:= a_{p, \bar{z}}^2 z_p^2 L_{\rho}^2 \Lambda \Gamma(1/a^*), \quad \text{for } p \geq 2. \end{aligned} \tag{3.11}$$

Clearly, $\hat{\gamma}_2 = \bar{\gamma}$.

Proposition 3.2 (*Upper bound on $\mathcal{K}(t, x)$*) *Suppose that $a \in]1, 2]$ and $|\delta| \leq 2 - a$. The kernel function $\mathcal{K}(t, x)$ defined in (2.9) satisfies, for all $t \geq 0$ and $x \in \mathbb{R}$,*

$$\mathcal{K}(t, x) \leq {}_{\delta} G_a(t, x) \frac{\gamma}{t^{1/a}} E_{1/a^*, 1/a^*} \left(\gamma t^{1/a^*} \right) \tag{3.12}$$

$$\leq \frac{C}{t^{1/a}} {}_{\delta} G_a(t, x) \left(1 + t^{1/a} \exp \left(\gamma^{a^*} t \right) \right), \tag{3.13}$$

where the constant $C = C(a, \delta, \lambda)$ can be chosen as

$$C(a, \delta, \lambda) := \gamma \sup_{t \geq 0} \frac{E_{1/a^*, 1/a^*} \left(\gamma t^{1/a^*} \right)}{1 + t^{1/a} \exp \left(\gamma^{a^*} t \right)} < +\infty. \tag{3.14}$$

This proposition is proved in Sect. 4. For a lower bound on $\mathcal{K}(t, x)$, we need another family of kernel functions: for $a > 0, t > 0$ and $x \in \mathbb{R}$, define

$$g_a(t, x) := \frac{1}{\pi} \frac{t}{(t^{2/a} + x^2)^{\frac{a}{2} + \frac{1}{2}}}. \tag{3.15}$$

These functions have the same scaling property as ${}_\delta G_a(t, x)$ (see Lemma 4.1(iv) below):

$$g_a(t, x) = \frac{1}{t^{1/a}} g_a\left(1, \frac{x}{t^{1/a}}\right).$$

Note that $g_1(t, x)$ is the Poisson kernel (see, e.g., [26, p. 268]), which satisfies the semigroup property

$$(g_1(t - s, \cdot) * g_1(s, \cdot))(x) = g_1(t, x), \quad 0 \leq s \leq t, x \in \mathbb{R}.$$

For $a \in]1, 2[$ and $|\delta| < 2 - a$, define

$$\tilde{C}_{a,\delta} := \inf_{(t,x) \in \mathbb{R}_+^* \times \mathbb{R}} \frac{{}_\delta G_a(t, x)}{\pi g_a(t, x)} > 0 \tag{3.16}$$

(for the strict positivity, see Lemma 5.1 below). Then let

$$\Upsilon(\lambda, a, \delta) := \frac{\lambda^2 \tilde{C}_{a,\delta}^2 C_{a+1/2}^2 \Gamma(1/a^*)}{\pi 2^{2(a+3+1/a)}}, \tag{3.17}$$

where

$$C_\nu := \frac{\Gamma(\nu)\Gamma(1/2)}{2\Gamma(\nu + 1/2)}, \quad \nu \geq 1/2. \tag{3.18}$$

Proposition 3.3 (Lower bound on $\mathcal{K}(t, x)$) *Fix $a \in]1, 2[$ and $|\delta| < 2 - a$ (note the strict inequality). Then for all $t > 0$ and $x \in \mathbb{R}$,*

$$\mathcal{K}(t, x) \geq \pi^2 \tilde{C}_{a,\delta}^2 \lambda^2 \Gamma(1/a^*) g_a^2(t, x) E_{1/a^*, 1/a^*} \left(\Upsilon(\lambda, a, \delta) t^{1/a^*} \right). \tag{3.19}$$

In particular, for all $t > 0$ and $x \in \mathbb{R}$,

$$(1 \star \mathcal{K})(t, x) \geq \frac{\pi^{1/2} \tilde{C}_{a,\delta}^2 \lambda^2 \Gamma(1/a^*) \Gamma(a + 1/2)}{\Gamma(1+a)} t^{1/a^*} E_{1/a^*, 1+1/a^*} \left(\Upsilon(\lambda, a, \delta) t^{1/a^*} \right). \tag{3.20}$$

This proposition is proved in Sect. 5.1.

3.3 Growth indices and weak intermittency

Theorem 3.4 (Weak intermittency) *Suppose that $a \in]1, 2]$ and $|\delta| \leq 2 - a$.*

(1) *If ρ satisfies (2.6) and $\mu \in \mathcal{M}_a(\mathbb{R})$, then for all even integers $p \geq 2$,*

$$\sup_{x \in \mathbb{R}} \bar{m}_p(x) \leq \frac{1}{2} \left(16 L_\rho^2 \Lambda \Gamma(1/a^*) \right)^{a^*} p^{2+1/(a-1)}. \tag{3.21}$$

(2) *Suppose ρ satisfies (1.3), $|\delta| < 2 - a$ (strict inequality) and $\mu \in \mathcal{M}_{a,+}(\mathbb{R})$. If either $\mu \neq 0$ or $\underline{\mu} \neq 0$, then for all $p \geq 2$,*

$$\inf_{x \in \mathbb{R}} \underline{m}_p(x) \geq \frac{p}{2} \Upsilon(l_\rho, a, \delta)^{a^*} > 0.$$

Note that if $a = 2$, then (3.21) implies that for some constant C , we have $\bar{m}_p \leq Cp^3$, which recovers previous analyses (see [3], [6, Example 2.7], etc).

Remark 3.5 Fix $p \geq 2$. Clearly, Theorem 3.4 implies that for all $x \in \mathbb{R}$,

$$\begin{aligned} \liminf_{t \rightarrow \infty} \frac{1}{t} \sup_{y \in \mathbb{R}} \log \mathbb{E}(|u(t, y)|^p) &\geq \liminf_{t \rightarrow \infty} \frac{1}{t} \log \mathbb{E}(|u(t, x)|^p) \\ &= \underline{m}_p(x) \geq \frac{p}{2} \Upsilon(l_\rho, a, \delta)^{a^*} > 0. \end{aligned}$$

Hence, the function $t \mapsto \sup_{y \in \mathbb{R}} \mathbb{E}(|u(t, y)|^p)$ has at least exponential growth. This answers the second open problem stated by Conus and Khoshnevisan in [10]. Moreover, Theorem 3.4 implies that for all fixed $x \in \mathbb{R}$, the function $t \mapsto \mathbb{E}(|u(t, x)|^p)$ also has at least exponential growth.

Recall the definitions of the constants $\widehat{\gamma}_p$ and $\Upsilon(l_\rho, a, \delta)$ in (3.11) and (3.17), respectively.

Theorem 3.6 (Growth indices)

(1) *Suppose that $a \in]1, 2]$, $|\delta| \leq 2 - a$ and ρ satisfies (2.6) with $\bar{\nu} = 0$. If there are $C < \infty$, $\alpha > 0$ and $\beta > 0$ such that for all $(t, x) \in [1, \infty[\times \mathbb{R}$,*

$$|J_0(t, x)| \leq C(1 + t^\alpha)(1 + |x|)^{-\beta}, \tag{3.22}$$

then

$$\bar{e}(p) \leq \frac{\widehat{\gamma}_p^{a^*}}{\beta} < +\infty. \tag{3.23}$$

In particular, if, for some $\eta > 0$, $\int_{\mathbb{R}} |\mu|(dy)(1 + |y|^\eta) < \infty$, then (3.22) and (3.23) are satisfied with $\beta = \min(\eta, 1 + a)$.

(2) Suppose that $a \in]1, 2[$ (note that $a \neq 2$), $|\delta| < 2 - a$ (strict inequality) and ρ satisfies (1.3). For all $\mu \in \mathcal{M}_{a,+}(\mathbb{R})$, $\mu \neq 0$ and all $p \geq 2$, if $\underline{\zeta} = 0$, then

$$\underline{e}(p) \geq \frac{\Upsilon(l_\rho, a, \delta)^{a^*}}{2(a+1)} > 0.$$

For these μ , if $\underline{\zeta} = 0$ and there is $c > 0$ such that

$$J_0(t, x) \geq c, \quad \text{for all } (t, x) \in \mathbb{R}_+ \times \mathbb{R}, \tag{3.24}$$

or if $\underline{\zeta} \neq 0$, then $\underline{e}(p) = \bar{e}(p) = +\infty$. In particular, $\underline{\lambda}(p) = \bar{\lambda}(p) = +\infty$ for all $p \geq 2$, and a sufficient condition for (3.24) is that $\mu(dx) = f(x)dx$ with $f(x) \geq c$, for all $x \in \mathbb{R}$.

The two theorems above are proved in Sect. 5.

Remark 3.7 In the case of the classical parabolic Anderson model, in which $a = 2$, $\delta = 0$ and $\rho(u) = \lambda u$, it was shown in [6] that $\underline{\lambda}(2) = \bar{\lambda}(2) = \lambda^2/2$ when the initial data has compact support (for instance). Here, it is natural to ask whether $\underline{e}(p) = \bar{e}(p)$ when $\rho(u) = \lambda u$, for instance for initial data with compact support. This remains an open question.

4 Proof of Theorem 3.1

We need some technical results. The proof of Theorem 3.1 will be presented at the end of this section.

The Green functions defined in (2.4) are densities of stable random variables. Some key properties are stated in the next lemma. Recall that a probability density function $f : \mathbb{R} \rightarrow \mathbb{R}_+$ is called *bell-shaped* if f is infinitely differentiable and its k -th derivative $f^{(k)}$ has exactly k zeros in its support for all k .

Lemma 4.1 *For $a \in]0, 2]$, the following properties hold:*

- (i) *For fixed $t > 0$, the function ${}_\delta G_a(t, \cdot)$ is a bell-shaped density function. In particular, $\int_{\mathbb{R}} {}_\delta G_a(t, x) dx = 1$.*
- (ii) *The unique mode is located on the positive semi-axis $x > 0$ if $\delta > 0$, on the negative semi-axis $x < 0$ if $\delta < 0$, and at $x = 0$ if $\delta = 0$.*
- (iii) *${}_\delta G_a(t, x)$ satisfies the semigroup property, i.e., for $0 < s < t$,*

$${}_\delta G_a(t + s, x) = ({}_\delta G_a(t, \circ) * {}_\delta G_a(s, \circ))(x).$$

- (iv) *${}_\delta G_a(t, x)$ has the following scaling property: For all $n \geq 0$,*

$$\frac{\partial^n}{\partial x^n} {}_\delta G_a(t, x) = t^{-\frac{n+1}{a}} \frac{\partial^n}{\partial \xi^n} {}_\delta G_a(1, \xi) \Big|_{\xi=t^{-1/a}x}. \tag{4.1}$$

(v) For $0 < a < 2$ with $a \neq 1$, when $x \rightarrow \pm\infty$,

$$\delta G_a(1, x) = \frac{1}{\pi} \sum_{j=1}^N |x|^{-aj-1} \frac{(-1)^{j+1}}{j!} \Gamma(aj + 1) \sin(j(a \pm \delta)\pi/2) + O(|x|^{-a(N+1)-1}).$$

(vi) If $a \in]1, 2]$, then there exist finite constants $K_{a,n}$ such that

$$\left| \frac{\partial^n}{\partial x^n} \delta G_a(1, x) \right| \leq \frac{K_{a,n}}{1 + |x|^{1+n+a}}, \quad \text{for } n \geq 0. \tag{4.2}$$

Moreover, for all $T \geq t > 0$, $n \geq 0$ and $x \in \mathbb{R}$,

$$\left| \frac{\partial^n}{\partial x^n} \delta G_a(t, x) \right| \leq t^{-\frac{n+1}{a}} \frac{K_{a,n}}{1 + |t^{-1/a}x|^{1+n+a}} \leq K_{a,n} t^{-\frac{n+1}{a}} \frac{(T \vee 1)^{1+\frac{n+1}{a}}}{1 + |x|^{1+n+a}}. \tag{4.3}$$

(vii) $\lim_{t \downarrow 0} \delta G_a(t, x) = \delta_0(x)$, where $\delta_0(x)$ is the Dirac delta function with unit mass at zero.

Proof Most of these properties appear in several books [18, 24, 27]. We refer the interested readers to [13, Lemma 1] for Properties (i) (except the bell-shaped density), (iii) and (iv). Formula (v) can be found in [18], (5.9.3) in Sects. 5.9 and (5.8.6) in Sect. 5.8; for $x \rightarrow -\infty$, use (5.8.2c) in this reference (note that the formula in [13, Lemma 1(vii)] is not quite correct for $x \rightarrow -\infty$). The proof that the density is bell-shaped is due to Gawronski [16]. Property (ii) can be found in the summary part of [27, Sect. 2.7, p. 143–147].

Now we prove (vi). Property (4.2) follows from [13, Corollary 1]. By the scaling property (4.1) and (4.2),

$$\left| \frac{\partial^n}{\partial x^n} \delta G_a(t, x) \right| \leq t^{-\frac{n+1}{a}} \frac{K_{a,n}}{1 + |t^{-1/a}x|^{1+n+a}} = t^{-\frac{n+1}{a}} \frac{K_{a,n} t^{1+\frac{n+1}{a}}}{t^{1+\frac{n+1}{a}} + |x|^{1+n+a}}.$$

Using the fact that the function $t \mapsto \frac{t}{t+z}$ is monotone increasing on \mathbb{R}_+ , the above quantity is less than

$$t^{-\frac{n+1}{a}} \frac{K_{a,n} (T \vee 1)^{1+\frac{n+1}{a}}}{(T \vee 1)^{1+\frac{n+1}{a}} + |x|^{1+n+a}} \leq t^{-\frac{n+1}{a}} \frac{K_{a,n} (T \vee 1)^{1+\frac{n+1}{a}}}{1 + |x|^{1+n+a}}.$$

This proves (4.3).

Property (vii) follows easily by taking Fourier transforms $\mathcal{F}(\delta G_a(t, \cdot))(\xi) = \exp(\delta \psi_a(\xi)t) \rightarrow 1$ as $t \downarrow 0$. This completes the proof of Lemma 4.1. \square

Let $\mathcal{L}_n(t, x; \lambda)$, $\mathcal{K}(t, x; \lambda)$, and $\Lambda = \delta \Lambda_a$ be defined in (2.8), (2.9), and (3.10), respectively. Recall that $1/a + 1/a^* = 1$.

Lemma 4.2 (Theorem 1.3 p. 32 in [22]) *If $0 < \alpha < 2$, β is an arbitrary complex number and μ is an arbitrary real number such that*

$$\pi\alpha/2 < \mu < \pi \wedge (\pi\alpha),$$

then for an arbitrary integer $p \geq 1$ the following expression holds:

$$E_{\alpha,\beta}(z) = \frac{1}{\alpha} z^{(1-\beta)/\alpha} \exp\left(z^{1/\alpha}\right) - \sum_{k=1}^p \frac{z^{-k}}{\Gamma(\beta - \alpha k)} + O\left(|z|^{-1-p}\right),$$

as $|z| \rightarrow \infty$ with $|\arg(z)| \leq \mu$.

Proposition 4.3 *For $1 < a \leq 2$, $|\delta| \leq 2 - a$ and $\lambda > 0$, we have the following properties:*

- (i) $\mathcal{L}_n(t, x; \lambda)$ is non-negative and for all $n \geq 0$ and $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$,

$$\mathcal{L}_n(t, x; \lambda) \leq B_{n+1}(t; \lambda) \delta G_a(t, x), \tag{4.4}$$

where

$$B_n(t; \lambda) := \lambda^{2n} \Lambda^n \frac{\Gamma(1/a^*)^n}{\Gamma(n/a^*)} t^{-1+n/a^*} \quad (n \geq 0, \lambda \in \mathbb{R}).$$

- (ii) $B_n(t; \lambda) \geq 0$ and for all $m \in \mathbb{N}^*$, $\sum_{n=0}^\infty B_n(t; \lambda)^{1/m} < +\infty$.
- (iii) For all $t > 0$ and $\lambda > 0$, the series $\sum_{n=1}^\infty \mathcal{L}_n(t, x; \lambda)$ converges uniformly over $x \in \mathbb{R}$ and hence $\mathcal{K}(t, x; \lambda)$ in (2.9) is well defined.

Proof (i) Non-negativity is clear. The scaling property (4.1) and the definition of Λ in (3.10) imply that

$$\delta G_a(t, x) \leq t^{-1/a} \Lambda, \tag{4.5}$$

which establishes the case $n = 0$ in (4.4). Suppose by induction that the relation (4.4) holds up to $n - 1$. Then by (4.5), we have

$$\begin{aligned} \mathcal{L}_n(t, x; \lambda) &= \int_0^t ds \int_{\mathbb{R}} dy \mathcal{L}_{n-1}(t-s, x-y) \lambda^2 \delta G_a^2(s, y) \\ &\leq \lambda^{2(n+1)} \Lambda^{n+1} \frac{\Gamma(1/a^*)^n}{\Gamma(n/a^*)} \int_0^t ds (t-s)^{-1+n/a^*} s^{-1/a} \\ &\quad \times \int_{\mathbb{R}} dy \delta G_a(t-s, x-y) \delta G_a(s, y). \end{aligned}$$

The conclusion now follows from the semigroup property of ${}_{\delta}G_a(t, x)$ and Euler's Beta integral (see [21, 5.12.1, on p. 142]):

$$\int_0^t ds s^{a-1} (t-s)^{b-1} = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)} t^{a+b-1}, \quad \text{with } \Re(a) > 0 \text{ and } \Re(b) > 0. \tag{4.6}$$

(ii) The non-negativity is clear. Denote $\beta := 1/a^*$. In order to show convergence of $\sum_n B_n(t; \lambda)^{1/m}$, we use the ratio test:

$$\left(\frac{B_n(t; \lambda)}{B_{n-1}(t; \lambda)} \right)^{1/m} = \left(\lambda^2 \Lambda \Gamma(\beta) t^\beta \right)^{1/m} \left(\frac{\Gamma((n-1)/a^*)}{\Gamma(n/a^*)} \right)^{1/m}.$$

By the asymptotic expansion of the Gamma function ([21, 5.11.2, on p. 140]),

$$\frac{\Gamma((n-1)/a^*)}{\Gamma(n/a^*)} \approx \left(\frac{e}{\beta} \right)^\beta \left(1 - \frac{1}{n} \right)^{(n-1)\beta} \frac{1}{n^\beta} \approx \frac{1}{(\beta n)^\beta},$$

for large n . Clearly, $\beta > 0$ since $1/a < 1$. Hence for all $t > 0$, for large n ,

$$\left(\frac{B_n(t; \lambda)}{B_{n-1}(t; \lambda)} \right)^{1/m} \approx \left(\lambda^2 \Lambda \Gamma(\beta) t^\beta \right)^{1/m} \frac{1}{(\beta n)^{\beta/m}},$$

and this goes to zero as $n \rightarrow +\infty$.

(iii) By (4.4) and (4.5),

$$\mathcal{L}_n(t, x; \lambda) \leq B_{n+1}(t; \lambda) t^{-1/a} \Lambda,$$

so (ii) implies (iii). This completes the proof of Proposition 4.3. □

Proof of Proposition 3.2 The bound (3.12) follows from the fact that

$$\sum_{k=1}^{\infty} \frac{z^k}{\Gamma(\alpha k)} = z E_{\alpha, \alpha}(z), \tag{4.7}$$

which can be easily seen from the definition, and the bound in Proposition 4.3(i):

$$\begin{aligned} \mathcal{K}(t, x; \lambda) &\leq {}_{\delta}G_a(t, x) \sum_{n=1}^{\infty} B_n(t; \lambda) = \frac{1}{t} {}_{\delta}G_a(t, x) \sum_{n=1}^{\infty} \frac{\left(\lambda^2 \Lambda \Gamma(1/a^*) t^{1/a^*} \right)^n}{\Gamma(n/a^*)} \\ &= \lambda^2 \Lambda \Gamma(1/a^*) t^{-1/a} {}_{\delta}G_a(t, x) E_{1/a^*, 1/a^*} \left(\lambda^2 \Lambda \Gamma(1/a^*) t^{1/a^*} \right). \end{aligned}$$

As for (3.13), we only need to show that the constant C defined in (3.14) is finite. Let

$$f(t) = \frac{E_{1/a^*, 1/a^*}(\gamma t^{1/a^*})}{1 + t^{1/a} \exp(\gamma^{a^*} t)}.$$

By Lemma 4.2 with the real non-negative value $z = \gamma t^{1/a^*}$ and $p = 1$,

$$\gamma E_{1/a^*, 1/a^*}(\gamma t^{1/a^*}) \leq a^* \gamma^{a^*} t^{1/a} \exp(\gamma^{a^*} t) + O\left(\frac{1}{|t|^{2/a^*}}\right), \quad t \rightarrow +\infty,$$

where we have used the convention that $1/\Gamma(0) = 0$ (see [21, 5.7.1, p.139]), therefore

$$\lim_{t \rightarrow +\infty} f(t) \leq a^* \gamma^{a^*}.$$

Since $E_{\alpha, \alpha}(\cdot)$ is continuous (by uniform convergence of the series in (3.9)), we conclude that $\sup_{t \geq 0} f(t) < +\infty$. This completes the proof of Proposition 3.2. \square

The next proposition is in principle a consequence of certain calculations in [13]. It is however not stated explicitly there, so we include a proof for the convenience of the reader.

Proposition 4.4 Fix $1 < a \leq 2$, $|\delta| \leq 2 - a$ and $1/a + 1/a^* = 1$. There are three universal constants

$$C_1 := \int_{\mathbb{R}} \frac{1 - \cos(u)}{2\pi \cos(\pi\delta/2)|u|^a} du, \quad C_3 := \frac{a^* \Gamma(1 + 1/a)}{2^{1/a} \pi \cos^{1/a}(\pi\delta/2)},$$

and

$$C_2 := (2^{1/a} - 1) C_3 + \frac{\Gamma(1 + 1/a) \sin^2(\pi\delta/2)}{2\pi a [\cos(\pi\delta/2)]^{2+1/a}},$$

such that

(i) for all $t > 0$ and $x, y \in \mathbb{R}$,

$$\int_0^t dr \int_{\mathbb{R}} dz [\delta G_a(t - r, x - z) - \delta G_a(t - r, y - z)]^2 \leq C_1 |x - y|^{a-1}; \tag{4.8}$$

(ii) for all $s, t \in \mathbb{R}_+^*$ with $s \leq t$, and $x \in \mathbb{R}$,

$$\int_0^s dr \int_{\mathbb{R}} dz [\delta G_a(t - r, x - z) - \delta G_a(s - r, x - z)]^2 \leq C_2 (t - s)^{1-1/a} \tag{4.9}$$

and

$$\int_s^t dr \int_{\mathbb{R}} dz [\delta G_a(t-r, x-z)]^2 \leq C_3(t-s)^{1-1/a}. \tag{4.10}$$

Remark 4.5 This proposition is a generalization of [6, Proposition 3.5] for the heat equation. In fact, if we take $a = 2$ and $\delta = 0$, then $\delta G_a(t, x) = G_2(t, x) = \frac{1}{\sqrt{4\pi t}} \exp(-x^2/(4t))$. Let $C'_i, i = 1, 2, 3$, be the optimal constants given in [6, Proposition 3.5] with $\nu = 2$. Then we have the following relations:

$$\frac{C'_1}{2} = C_1 = \frac{1}{2}, \quad \frac{C'_2}{\sqrt{2}} = C_2 = \frac{\sqrt{2}-1}{\sqrt{2\pi}}, \quad \frac{C'_3}{\sqrt{2}} = C_3 = \frac{1}{\sqrt{2\pi}},$$

where for C_1 , we use the fact that $\int_{\mathbb{R}} \frac{1-\cos(u)}{u^2} du = 2 \int_0^\infty \frac{\sin(u)}{u} du = \pi$; see [21, 4.26.12, on p. 122] for the last integral. This recovers the optimal inequalities of [6, Proposition 3.5], since ν appears in the right-hand side of the inequalities in this reference.

Proof of Proposition 4.4 (i) Note that

$$\begin{aligned} \mathcal{F}(\delta G_a(t, \cdot))(\xi) &:= \int_{\mathbb{R}} dx e^{-i\xi x} \delta G_a(t, x) = \exp\{t \delta \psi_a(\xi)\} \\ &= \exp\left\{-t|\xi|^a e^{-i\pi\delta \operatorname{sgn}(\xi)/2}\right\}. \end{aligned}$$

By Plancherel’s theorem, the left hand side of (4.8) equals

$$\begin{aligned} &\frac{1}{2\pi} \int_0^t dr \int_{\mathbb{R}} d\xi \left| e^{-i\xi x - (t-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2}} - e^{-i\xi y - (t-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2}} \right|^2 \\ &= \frac{1}{2\pi} \int_0^t dr \int_{\mathbb{R}} d\xi \exp\{-2(t-r)|\xi|^a \cos(\pi\delta/2)\} \left| e^{-i\xi x} - e^{-i\xi y} \right|^2 \\ &= \frac{1}{\pi} \int_0^t dr \int_{\mathbb{R}} d\xi \exp\{-2(t-r)|\xi|^a \cos(\pi\delta/2)\} (1 - \cos(\xi(x-y))). \end{aligned}$$

After integrating over r , the above integral equals

$$\frac{1}{\pi} \int_{\mathbb{R}} d\xi \frac{1 - \exp\{-2t|\xi|^a \cos(\pi\delta/2)\}}{2 \cos(\pi\delta/2)|\xi|^a} (1 - \cos(\xi(x-y))).$$

Use the change of variables $\xi = u/|x-y|$ to see that this is equal to

$$\begin{aligned} &\frac{1}{\pi} |x-y|^{a-1} \int_{\mathbb{R}} du \frac{1 - \exp(-2t|u|^a \cos(\pi\delta/2)/|x-y|^a)}{2 \cos(\pi\delta/2)|u|^a} (1 - \cos(u)) \\ &\leq C'_1 |x-y|^{a-1}, \end{aligned}$$

where

$$C'_1 = \int_{\mathbb{R}} \frac{1 - \cos(u)}{2\pi \cos(\pi\delta/2)|u|^a} du.$$

This proves (4.8).

(ii) Denote the left hand side of (4.9) by I . Apply Plancherel's theorem to I :

$$\begin{aligned} I &= \frac{1}{2\pi} \int_0^s dr \int_{\mathbb{R}} d\xi |\exp(-i\xi x - (t-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2}) \\ &\quad - \exp(-i\xi x - (s-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2})|^2 \\ &= \frac{1}{2\pi} \int_0^s dr \int_{\mathbb{R}} d\xi |\exp(-(t-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2}) \\ &\quad - \exp(-(s-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2})|^2. \end{aligned}$$

Denote $\beta := \pi\delta \operatorname{sgn}(\xi)/2 \in]-\frac{\pi}{2}, \frac{\pi}{2}[$ and

$$A_{r,t} := (t-r)|\xi|^a \cos(\beta), \quad B_{r,t} := (t-r)|\xi|^a \sin(\beta).$$

Then

$$\begin{aligned} &\left| \exp(-(t-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2}) - \exp(-(s-r)|\xi|^a e^{i\pi\delta \operatorname{sgn}(\xi)/2}) \right|^2 \\ &= \left| e^{-A_{r,t}} \cos(B_{r,t}) - i e^{-A_{r,t}} \sin(B_{r,t}) - e^{-A_{r,s}} \cos(B_{r,s}) + i e^{-A_{r,s}} \sin(B_{r,s}) \right|^2 \\ &= e^{-2A_{r,t}} + e^{-2A_{r,s}} - 2e^{-(A_{r,t}+A_{r,s})} \cos(B_{r,t} - B_{r,s}). \end{aligned}$$

Now, according to [21, Equation 5.9.1, p.139], for $z \in \mathbb{C}$ with $\Re(z) > 0$,

$$\int_{\mathbb{R}} d\xi e^{-z|\xi|^a} = 2 \int_0^\infty d\xi e^{-z\xi^a} = 2z^{-1/a} \Gamma(1 + 1/a). \tag{4.11}$$

Hence, since $\cos(\beta) = \cos(\pi\delta/2)$, which does not depend on ξ ,

$$\int_{\mathbb{R}} d\xi e^{-2A_{r,t}} = \int_{\mathbb{R}} d\xi e^{-2(t-r)\cos(\beta)|\xi|^a} = \frac{2^{1/a} \Gamma(1 + 1/a)}{\cos^{1/a}(\beta)} \frac{1}{(t-r)^{1/a}}. \tag{4.12}$$

Similarly,

$$\int_{\mathbb{R}} d\xi e^{-2A_{r,s}} = \frac{2^{1/a} \Gamma(1 + 1/a)}{\cos^{1/a}(\beta)} \frac{1}{(s-r)^{1/a}}.$$

For the third term, notice that

$$\begin{aligned} &e^{-(A_{r,t}+A_{r,s})} \cos(B_{r,t} - B_{r,s}) \\ &= \exp\left(-\left(\frac{t+s}{2} - r\right) 2|\xi|^a \cos(\beta)\right) \cos((t-s)|\xi|^a \sin(\beta)) \end{aligned}$$

$$= \Re \left[\exp \left\{ - \left[\left(\frac{t+s}{2} - r \right) 2 \cos(\beta) + i(t-s) \sin(\beta) \right] |\xi|^a \right\} \right].$$

Apply (4.11) with $z = \left(\frac{t+s}{2} - r \right) 2 \cos(\beta) + i(t-s) \sin(\beta)$:

$$\begin{aligned} & \int_{\mathbb{R}} d\xi \exp \left\{ - \left[\left(\frac{t+s}{2} - r \right) 2 \cos(\beta) + i(t-s) \sin(\beta) \right] |\xi|^a \right\} \\ &= 2\Gamma(1 + 1/a) \left[\left(\frac{t+s}{2} - r \right) 2 \cos(\beta) + i(t-s) \sin(\beta) \right]^{-1/a}. \end{aligned}$$

Apply Lemma 4.6 below with $c = 1/a$, $b = \left(\frac{t+s}{2} - r \right) 2 \cos(\beta)$ and $x = (t-s)^2 \sin^2(\beta)$:

$$\begin{aligned} & \Re \left[\left(\frac{t+s}{2} - r \right) 2 \cos(\beta) + i(t-s) \sin(\beta) \right]^{-1/a} \\ & \geq \frac{1}{2^{1/a} \cos^{1/a}(\beta)} \frac{1}{((t+s)/2 - r)^{1/a}} \\ & \quad - \frac{(a+1) \sin^2(\beta)}{2a^2 [2 \cos(\beta)]^{2+1/a}} \frac{(t-s)^2}{((t+s)/2 - r)^{2+1/a}}. \end{aligned}$$

Hence,

$$\begin{aligned} & 2 \int_{\mathbb{R}} d\xi e^{-(A_{r,t} + A_{r,s})} \cos(B_{r,t} - B_{r,s}) \\ & \geq 2^{1+1/a^*} \frac{\Gamma(1 + 1/a)}{\cos^{1/a}(\beta)} \frac{1}{((t+s)/2 - r)^{1/a}} \\ & \quad - \frac{2\Gamma(1 + 1/a)(a+1) \sin^2(\beta)}{a^2 [2 \cos(\beta)]^{2+1/a}} \frac{(t-s)^2}{((t+s)/2 - r)^{2+1/a}}. \end{aligned}$$

Integrating over r and then applying Lemma 4.7 below, we get

$$\begin{aligned} I & \leq \frac{\Gamma(1 + 1/a)}{2^{1/a} \pi \cos^{1/a}(\beta)} \int_0^s dr \left(\frac{1}{(t-r)^{1/a}} + \frac{1}{(s-r)^{1/a}} - \frac{2}{[(t+s)/2 - r]^{1/a}} \right) \\ & \quad + \frac{\Gamma(1 + 1/a)(a+1) \sin^2(\beta)}{\pi a^2 [2 \cos(\beta)]^{2+1/a}} \int_0^s dr \frac{(t-s)^2}{((t+s)/2 - r)^{2+1/a}} \\ & \leq C_2(t-s)^{1/a^*}, \end{aligned}$$

where $1/a^* + 1/a = 1$.

As for (4.10), from (4.12), we have

$$\begin{aligned} \int_s^t dr \int_{\mathbb{R}} dz [\delta G_a(t-r, x-z)]^2 &= \frac{1}{2\pi} \int_s^t dr \int_{\mathbb{R}} d\xi e^{-2(t-r)|\xi|^a \cos(\beta)} \\ &= \frac{\Gamma(1 + 1/a)}{2^{1/a} \pi \cos^{1/a}(\beta)} \int_s^t dr \frac{1}{(t-r)^{1/a}} \\ &= \frac{a^* \Gamma(1 + 1/a)}{2^{1/a} \pi \cos^{1/a}(\beta)} (t-s)^{1/a^*}. \end{aligned} \tag{4.13}$$

This completes the proof of Proposition 4.4. □

Lemma 4.6 *Suppose $b > 0$ and $c \in]0, 1]$. Then for all $x \geq 0$,*

$$\Re \left((b \pm i\sqrt{x})^{-c} \right) \geq \frac{1}{b^c} - \frac{c(1+c)}{2} \frac{x}{b^{2+c}}.$$

Proof Let $\theta = \arctan(\sqrt{x}/b) \in [0, \pi/2[$ and denote $f(x) := \Re \left((b \pm i\sqrt{x})^{-c} \right)$. Then

$$f(x) = \left(b^2 + x \right)^{-c/2} \cos(c\theta).$$

Because $\cos(\theta) \geq 1 - \theta^2/2$ and $\arctan(y) \leq y$ for $y \geq 0$, we have that

$$\cos(c\theta) \geq 1 - \frac{c^2\theta^2}{2} \geq 1 - \frac{c^2x}{2b^2}.$$

By Taylor's theorem, for some $\zeta \in [0, x]$,

$$\left(b^2 + x \right)^{-c/2} = b^{-c} - \frac{1}{2}cx(b^2 + \zeta)^{-1-c/2} \geq b^{-c} - \frac{1}{2}cxb^{-2-c}.$$

Combining the above two lower bounds proves the lemma. □

Lemma 4.7 *For all $t \geq s \geq 0$ and $a \in]1, 2]$, we have*

$$\int_0^s dr \left(\frac{1}{(t-r)^{1/a}} + \frac{1}{(s-r)^{1/a}} - \frac{2}{((t+s)/2-r)^{1/a}} \right) \leq a^*(2^{1/a}-1) (t-s)^{1/a^*},$$

and

$$\int_0^s dr \frac{(t-s)^2}{((t+s)/2-r)^{2+1/a}} \leq \frac{a}{a+1} 2^{1+1/a} (t-s)^{1/a^*},$$

where a^* is the dual of a : $1/a + 1/a^* = 1$.

Proof Clearly,

$$\begin{aligned} & \frac{1}{a^*} \int_0^s dr \left(\frac{1}{(t-r)^{1/a}} + \frac{1}{(s-r)^{1/a}} - \frac{2}{((t+s)/2-r)^{1/a}} \right) \\ &= s^{1/a^*} + t^{1/a^*} - (t-s)^{1/a^*} + 2^{1/a}(t-s)^{1/a^*} - 2^{1/a}(t+s)^{1/a^*}. \end{aligned}$$

We shall to prove that

$$(t-s)^{-1/a^*} [s^{1/a^*} + t^{1/a^*} - (t-s)^{1/a^*} + 2^{1/a}(t-s)^{1/a^*} - 2^{1/a}(t+s)^{1/a^*}]$$

is bounded from above for all $0 \leq s < t$, or, equivalently, that

$$g(r) := \frac{r^{1/a^*} + 1 - (1-r)^{1/a^*} + 2^{1/a}(1-r)^{1/a^*} - 2^{1/a}(1+r)^{1/a^*}}{(1-r)^{1/a^*}}$$

is bounded for all $r \in [0, 1[$. Clearly, $g(0) = 0$ and $\lim_{r \uparrow 1} g(r) = 2^{1/a} - 1$ (by applying L'Hospital's rule once). Hence $\sup_{r \in [0, 1[} g(r) < \infty$. In addition,

$$g'(r) = \frac{((1+r)^{1/a} + (1+1/r)^{1/a}) - 2^{1+1/a}}{a^*(1-r)^{2-1/a}(1+r)^{1/a}},$$

and notice that for all $r \in]0, 1[$,

$$(1+r)^{1/a} + (1+1/r)^{1/a} \geq 2[(1+r)(1+1/r)]^{1/(2a)} = 2\left(\sqrt{r} + \frac{1}{\sqrt{r}}\right)^{1/a} \geq 2^{1+1/a}.$$

Hence $g'(r) \geq 0$ for $r \in [0, 1[$ and $\sup_{r \in [0, 1[} g(r) = \lim_{r \uparrow 1} g(r) = 2^{1/a} - 1$. Therefore, the first inequality is proved with the constant $a^*(2^{1/a} - 1)$.

As for the second inequality, we have that

$$\begin{aligned} \int_0^s dr \frac{(t-s)^2}{((t+s)/2-r)^{2+1/a}} &= \frac{a}{a+1} 2^{1+1/a} \frac{((t+s)^{1+1/a} - (t-s)^{1+1/a})}{(t+s)^{1+1/a}} (t-s)^{1/a^*} \\ &\leq \frac{a}{a+1} 2^{1+1/a} (t-s)^{1/a^*}, \end{aligned}$$

which completes the proof of the Lemma 4.7. □

The following proposition is useful to prove the $L^p(\Omega)$ -continuity of $I(t, x)$.

Proposition 4.8 *Suppose that $a \in]1, 2]$ and $|\delta| \leq 2 - a$. Fix $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$. Denote*

$$B := B_{t,x} = \{(t', x') \in \mathbb{R}_+^* \times \mathbb{R} : 0 \leq t' \leq t + 1/2, |x - x'| \leq 1\}.$$

Then there exists a constant $A > 0$ such that for all $(t', x') \in B$, $s \in [0, t'[$ and $|y| \geq A$,

$$\delta G_a(t' - s, x' - y) \leq -\delta G_a(t + 1 - s, x - y) + \delta G_a(t + 1 - s, x - y).$$

Proof The case where $a = 2$ is proved in [6, Proposition A.3], so we only need to prove the case where $1 < a < 2$. Denote $F(t, x) := \delta G_a(t, x) + -\delta G_a(t, x)$. Suppose the mode of the density $\delta G_a(1, x)$ is located at $m \in \mathbb{R}$. By the scaling property, the mode of the density $\delta G_a(t, x)$ is located at $t^{1/a}m$. Hence, when $x \geq t^{1/a}|m|$ (resp. $x \leq -t^{1/a}|m|$), the function $x \mapsto F(t, x)$ is decreasing (resp. increasing).

Fix $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$. Assume that $|y - x| > 1 + (t + 1/2)^{1/a} |m|$. From the above observation, we deduce that for all $(t', x') \in B$,

$$\delta G_a(t' - s, x' - y) \leq F(t' - s, |y - x| - |x - x'|) \leq F(t' - s, |y - x| - 1). \tag{4.14}$$

Apply Lemma 4.1(v) with $N = 1$ and use the scaling property of $\delta G_a(t, x)$ to get

$$F(t, x) = 2 \frac{\Gamma(a + 1)}{\pi} \sin\left(\frac{\pi}{2} a\right) \cos\left(\frac{\pi}{2} \delta\right) t |x|^{-1-a} + O\left(t^2 |x|^{-1-2a}\right).$$

Because $|\delta| \leq 2 - a$ and $a \in]1, 2[$, we see that $\sin(\pi a/2) \cos(\pi \delta/2) \neq 0$. Hence,

$$\begin{aligned} & \frac{F(t + 1 - s, x - y)}{F(t' - s, |y - x| - 1)} \\ &= \frac{(t + 1 - s) |x - y|^{-1-a} + O\left((t + 1 - s)^2 |x - y|^{-1-2a}\right)}{(t' - s) \left(|y - x| - 1\right)^{-1-a} + O\left((t' - s)^2 \left(|y - x| - 1\right)^{-1-2a}\right)} \\ &= \frac{t + 1 - s}{t' - s} \frac{|x - y|^{-1-a} + O\left((t + 1 - s) |x - y|^{-1-2a}\right)}{\left(|y - x| - 1\right)^{-1-a} + O\left((t' - s) \left(|y - x| - 1\right)^{-1-2a}\right)}. \end{aligned}$$

Now it is clear that

$$\lim_{|y| \rightarrow +\infty} \inf_{(t', x') \in B, s \in [0, t'[} \frac{|x - y|^{-1-a} + O\left((t + 1 - s) |x - y|^{-1-2a}\right)}{\left(|y - x| - 1\right)^{-1-a} + O\left((t' - s) \left(|y - x| - 1\right)^{-1-2a}\right)} = 1,$$

which implies that

$$\begin{aligned} \lim_{|y| \rightarrow +\infty} \inf_{(t', x') \in B, s \in [0, t'[} \frac{F(t + 1 - s, x - y)}{F(t' - s, |y - x| - 1)} &\geq \inf_{(t', x') \in B, s \in [0, t'[} \frac{t + 1 - s}{t + 1/2 - s} \\ &= \frac{t + 1}{t + 1/2} = 1 + \frac{1}{2t + 1} > 1, \end{aligned}$$

where we have used the fact that $s \mapsto (t + 1 - s)/(t + 1/2 - s)$ is increasing. Hence, by (4.14), we can choose a large constant A such that for all $|y| \geq A$, $(t', x') \in B$ and $s \in [0, t'[$, the inequality

$$\frac{F(t + 1 - s, x - y)}{\delta G_a(t' - s, x' - y)} \geq 1 + \frac{1}{2(t + 1)} > 1$$

holds. This completes the proof of Proposition 4.8. □

Lemma 4.9 *For all $m, n \in \mathbb{N}$, there exist polynomials $\{P_i^{(n,m)}(x), i = 0, \dots, n\}$, such that:*

(1) the $P_i^{(n,m)}(x)$ are of degree $\leq i$ and they satisfy

$$\frac{\partial^{n+m}}{\partial t^n \partial x^m} \delta G_a(t, x) = \frac{1}{(at)^n} \sum_{i=0}^n P_i^{(n,m)}(x) \frac{\partial^{i+m}}{\partial x^{i+m}} \delta G_a(t, x);$$

(2) For fixed $t > 0$, the partial derivative $\frac{\partial^{n+m}}{\partial t^n \partial x^m} \delta G_a(t, \cdot)$ as a function of x is smooth and integrable.

Proof Part (2) is a direct consequence of (1) and (i) and (vi) in Lemma 4.1. We now prove (1). It is clearly true for $n = m = 0$: in this case, $P_0^{(0,0)}(x) \equiv 1$. Moreover, if $n = 0$, then it is trivially true, with $P_0^{(0,m)}(x) = 1$. Consider the case $n = 1$ and $m = 0$. Using the scaling properties twice, we have

$$\begin{aligned} \frac{\partial}{\partial t} \delta G_a(t, x) &= \left[-\frac{1/a}{t^{1+1/a}} \delta G_a(1, \xi) - \frac{1}{t^{1/a}} \frac{\partial \delta G_a(1, \xi)}{\partial \xi} \frac{x/a}{t^{1+1/a}} \right] \Big|_{\xi=t^{-1/a}x} \\ &= -\frac{1}{at} \left(\frac{1}{t^{1/a}} \delta G_a\left(1, \frac{x}{t^{1/a}}\right) + \frac{x}{t^{2/a}} \frac{\partial \delta G_a(1, \xi)}{\partial \xi} \Big|_{\xi=t^{-1/a}x} \right) \\ &= -\frac{1}{at} \left(\delta G_a(t, x) + x \frac{\partial}{\partial x} \delta G_a(t, x) \right). \end{aligned}$$

So in this case, $P_0^{(1,0)}(x) = -1$ and $P_1^{(1,0)}(x) = -x$. Now suppose that (1) is true for $n, m \in \mathbb{N}$. It is easy to see that (1) is true also for $n, m + 1$ with

$$\begin{aligned} P_i^{(n,m+1)}(x) &= P_i^{(n,m)}(x) + \frac{d}{dx} P_{i+1}^{(n,m)}(x), \quad \text{for } i = 0, \dots, n-1, \\ P_n^{(n,m+1)}(x) &= P_n^{(n,m)}(x), \end{aligned}$$

so $P_i^{(n,m+1)}(x)$ is a polynomial of degree $\leq i$.

Now assume by induction that $n \geq 1$ and the property is true for $\tilde{n} \leq n$ and all $m \geq 0$. We shall establish the property for $n + 1$ and m . By the induction assumption, we have

$$\begin{aligned} \frac{\partial^{n+1+m}}{\partial t^{n+1} \partial x^m} \delta G_a(t, x) &= \frac{-na}{(at)^{n+1}} \sum_{i=0}^n P_i^{(n,m)}(x) \frac{\partial^{i+m}}{\partial x^{i+m}} \delta G_a(t, x) \\ &\quad + \frac{1}{(at)^n} \sum_{i=0}^n P_i^{(n,m)}(x) \frac{\partial^{1+i+m}}{\partial t \partial x^{i+m}} \delta G_a(t, x). \end{aligned}$$

Then replace $\frac{\partial^{1+i+m}}{\partial t \partial x^{i+m}} \delta G_a(t, x)$ by the following sum using the induction assumption:

$$\frac{\partial^{1+i+m}}{\partial t \partial x^{i+m}} \delta G_a(t, x) = \frac{1}{at} \left(P_0^{(1,i+m)}(x) \frac{\partial^{i+m}}{\partial x^{i+m}} \delta G_a(t, x) \right)$$

$$+ P_1^{(1,i+m)}(x) \frac{\partial^{i+m+1}}{\partial x^{i+m+1}} \delta G_a(t, x) \Big).$$

Finally, after grouping terms one can choose the following polynomials:

$$P_0^{(n+1,m)}(x) = -na P_0^{(n,m)}(x) + P_0^{(n,m)}(x) P_0^{(1,m)}(x),$$

which is a polynomial of order 0,

$$P_i^{(n+1,m)}(x) = -na P_i^{(n,m)}(x) + P_i^{(n,m)}(x) P_0^{(1,i+m)}(x) + P_{i-1}^{(n,m)}(x) P_1^{(1,i+m-1)}(x),$$

which are polynomials of degree $\leq i$, for $i = 1, \dots, n$, and

$$P_{n+1}^{(n+1,m)}(x) = P_n^{(n,m)}(x) P_1^{(1,n+m)}(x),$$

which are polynomials of degree $\leq n + 1$. This completes the proof of Lemma 4.9. \square

Lemma 4.10 *Suppose that $a \in]1, 2]$ and $\mu \in \mathcal{M}_a(\mathbb{R})$.*

- (1) *The function $J_0(t, x) = (\delta G_a(t, \cdot) * \mu)(x)$ belongs to $C^\infty(\mathbb{R}_+^* \times \mathbb{R})$.*
- (2) *For all compact sets $K \subset \mathbb{R}_+^* \times \mathbb{R}$ and $v \in \mathbb{R}$,*

$$\sup_{(t,x) \in K} \left([v^2 + J_0^2] \star \mathcal{K} \right) (t, x) < \infty. \tag{4.15}$$

In fact, for all $(t, x) \in \mathbb{R}_+^ \times \mathbb{R}$,*

$$\left(J_0^2 \star \mathcal{K} \right) (t, x) \leq C'(t \vee 1)^{2(1+1/a)} t^{1-2/a} \left[t^{-1/a} + \exp(\gamma a^* t) \right], \tag{4.16}$$

where

$$C' := C A_a^2 K_{a,0}^2 \max \left(a^*, \frac{\Gamma(1/a^*)^2}{\Gamma(2/a^*)} \right), \tag{4.17}$$

$C = C(a, \delta, \lambda)$ is defined in (3.14), $K_{a,0}$ is defined in (4.2), and

$$A_a := \sup_{y \in \mathbb{R}} \int_{\mathbb{R}} \frac{|\mu|(dz)}{1 + |y - z|^{1+a}}. \tag{4.18}$$

Proof (1) Fix $0 < t \leq T$ and $n, m \in \mathbb{N}$. By Lemma 4.9 and (4.3),

$$\left| \frac{\partial^{n+m}}{\partial t^n \partial x^m} \delta G_a(t, x) \right| \leq \frac{1}{(at)^n} \sum_{i=0}^n \left| P_i^{(n,m)}(x) \right| K_{a,i+m} t^{-(i+m+1)/a} \frac{(T \vee 1)^{1+(i+m+1)/a}}{1 + |x|^{1+i+m+a}}.$$

Since the polynomials $P_i^{(n,m)}(x)$ are of degree $\leq i$, for some finite constant $C > 0$ depending on a, m, n and T , the above bound reduces to

$$\left| \frac{\partial^{n+m}}{\partial t^n \partial x^m} \delta G_a(t, x) \right| \leq C \frac{g(t)}{1 + |x|^{m+1+a}}, \quad \text{with } g(t) := \sum_{i=0}^n t^{-n-(i+m+1)/a}.$$

Hence, for $0 < t_1 < t_2 \leq T$,

$$\int_{t_1}^{t_2} ds \int_{\mathbb{R}} \mu(dz) \left| \frac{\partial^{n+m}}{\partial s^n \partial x^m} \delta G_a(s, z) \right| < +\infty. \tag{4.19}$$

By Fubini's theorem and induction, it is now possible to conclude that $J_0(\cdot, \circ) \in C^\infty(\mathbb{R}_+^* \times \mathbb{R})$. Indeed, the first step of this induction argument is:

$$\begin{aligned} J_0(t_2, x) - J_0(t_1, x) &= \int_{\mathbb{R}} \mu(dy) (\delta G_a(t_2, x - y) - \delta G_a(t_1, x - y)) \\ &= \int_{\mathbb{R}} \mu(dy) \int_{t_1}^{t_2} dt \frac{\partial}{\partial t} \delta G_a(t, x - y) \\ &= \int_{t_1}^{t_2} dt \int_{\mathbb{R}} \mu(dy) \frac{\partial}{\partial t} \delta G_a(t, x - y), \end{aligned}$$

where we have used Fubini's theorem, which applies by (4.19). This shows that

$$\frac{\partial}{\partial t} J_0(t, x) = \int_{\mathbb{R}} \mu(dy) \frac{\partial}{\partial t} \delta G_a(t, x - y),$$

and higher derivatives are obtained by induction. This proves (1).

(2) By (4.3), for $0 < s \leq t$,

$$|J_0(s, y)| \leq A_a K_{a,0} (t \vee 1)^{1+1/a} s^{-1/a}, \tag{4.20}$$

where A_a is defined in (4.18). Let $(t, x) \in \mathbb{R}_+^* \times \mathbb{R}$. By (3.13), and by replacing one factor $|J_0(s, y)|$ of $J_0^2(s, y)$ by the above bound, we have that

$$\begin{aligned} (J_0^2 \star \mathcal{K})(t, x) &\leq C \int_0^t ds \left(\frac{1}{(t-s)^{1/a}} + \exp(\gamma^{a^*}(t-s)) \right) \int_{\mathbb{R}} dy \delta G_a(t-s, x-y) \\ &\quad \times A_a K_{a,0} (t \vee 1)^{1+1/a} s^{-1/a} \left| \int_{\mathbb{R}} \mu(dz) \delta G_a(s, y-z) \right|, \end{aligned}$$

where the constant $C := C(a, \delta, \lambda)$ is defined in (3.14). Integrate over dy using the semigroup property, and then integrate over $\mu(dz)$:

$$(J_0^2 \star \mathcal{K})(t, x) \leq C A_a K_{a,0} (t \vee 1)^{1+1/a} |J_0(t, x)|$$

$$\times \int_0^t ds \frac{1}{s^{1/a}} \left[\frac{1}{(t-s)^{1/a}} + \exp\left(\gamma^{a^*}(t-s)\right) \right]. \tag{4.21}$$

Apply (4.20) to $J_0(t, x)$. The integral over s gives

$$\begin{aligned} & \int_0^t ds \left[\frac{1}{s^{1/a}(t-s)^{1/a}} + \frac{1}{s^{1/a}} \exp\left(\gamma^{a^*}(t-s)\right) \right] \\ & \leq \int_0^t ds \left[\frac{1}{s^{1/a}(t-s)^{1/a}} + \frac{1}{s^{1/a}} \exp\left(\gamma^{a^*}t\right) \right]. \end{aligned}$$

Use (4.6) to see that this is equal to

$$t^{1-2/a} \frac{\Gamma(1-1/a)^2}{\Gamma(2-2/a)} + a^* t^{1/a^*} \exp\left(\gamma^{a^*}t\right) = t^{1/a^*} \left[\frac{1}{t^{1/a}} \frac{\Gamma(1/a^*)^2}{\Gamma(2/a^*)} + a^* \exp\left(\gamma^{a^*}t\right) \right]. \tag{4.22}$$

Hence, combining the above facts proves (4.16).

For (4.15), we consider the case $\mu(dx) = v dx$, for which $\mu \in \mathcal{M}_a(\mathbb{R})$ and $J_0(t, x) \equiv v$. Together with (4.16) for all $\mu \in \mathcal{M}_a(\mathbb{R})$, we obtain (4.15). This completes the proof of Lemma 4.10. \square

Proof of Theorem 3.1 The proof follows the same six steps as those in the proof of [6, Theorem 2.4] with some minor changes:

- (1) Both proofs rely on estimates on the kernel function $\mathcal{K}(t, x)$. Instead of an explicit formula as for the heat equation case (see [6, Proposition 2.2]), Proposition 3.2 ensures the finiteness and provides a bound on the kernel function $\mathcal{K}(t, x)$.
- (2) In the Picard iteration scheme (i.e., Steps 1–4 in the proof of [6, Theorem 2.4]), we need to check the $L^p(\Omega)$ -continuity of the stochastic integral, which then guarantees that at the next step, the integrand is again in \mathcal{P}_2 , via [6, Proposition 3.4]. Here, the statement of [6, Proposition 3.4] is still true by replacing in its proof [6, Propositions 3.5 and A.3] by Propositions 4.4 and 4.8, respectively. Note that when applying Proposition 4.8, we need to replace the G_v^2 in [6, (3.8)] by $(-\delta G_a + \delta G_a)^2 \leq 2-\delta G_a^2 + 2\delta G_a^2$.
- (3) In the first step of the Picard iteration scheme, the following property is useful: For all compact sets $K \subseteq \mathbb{R}_+ \times \mathbb{R}$,

$$\sup_{(t,x) \in K} \left(\left[1 + J_0^2 \right] \star \delta G_a^2 \right) (t, x) < +\infty.$$

For the heat equation, this property is discussed in [6, Lemma 3.9]. Here, Lemma 4.10 gives the desired result with minimal requirements on the initial data. This property, together with the calculation of the upper bound on the function \mathcal{K} in Proposition 3.2, guarantees (as in [6, Lemmas 3.3 and 3.7]) that all the $L^p(\Omega)$ -moments of $u(t, x)$ are finite. This property is also used to establish uniform convergence of the Picard iteration scheme, hence $L^p(\Omega)$ -continuity of $(t, x) \mapsto I(t, x)$.

The proofs of (3.1)–(3.4) are identical to those of the corresponding properties in [6, Theorem 2.4], and (3.5) and (3.6) are direct consequences of the preceding statements.

This completes the proof of Theorem 3.1. □

5 Proofs of Theorems 3.4 and 3.6

We begin with the upper bound in Theorem 3.4.

Proof of Theorem 3.4(1) Recall from (3.11) that $\widehat{\gamma}_p = a_{p,\bar{\zeta}}^2 z_p^2 L_\rho^2 \Lambda \Gamma(1/a^*)$, and $a^* = a/(a - 1)$. By (3.1), (4.16) and (4.20), for all $x \in \mathbb{R}$,

$$\bar{m}_p(x) = \limsup_{t \rightarrow \infty} \frac{\log \|u(t, x)\|_p^p}{t} \leq \frac{\widehat{\gamma}_p^{a^*} P}{2} = \frac{P}{2} \left[a_{p,\bar{\zeta}}^2 z_p^2 L_\rho^2 \Lambda \Gamma(1/a^*) \right]^{a^*}.$$

Since $a_{p,\bar{\zeta}} \leq 2$ and $z_p \leq 2\sqrt{p}$, (3.21) follows. □

5.1 Lower bound on $\mathcal{K}(t, x)$ (Proposition 3.3)

We need some properties of $g_a(t, x)$ defined in (3.15).

Lemma 5.1 *Suppose that $a \in]1, 2[$ and $|\delta| < 2 - a$. Then the constant $\widetilde{C}_{a,\delta}$ defined in (3.16) is strictly positive, and so for all $t > 0$ and $x \in \mathbb{R}$,*

$$\delta G_a(t, x) \geq \widetilde{C}_{a,\delta} \pi g_a(t, x) = \frac{\widetilde{C}_{a,\delta} t}{(t^{2/a} + x^2)^{\frac{a}{2} + \frac{1}{2}}}. \tag{5.1}$$

Proof By the scaling property of both δG_a and $g_a(t, x)$,

$$\inf_{(t,x) \in \mathbb{R}_+^* \times \mathbb{R}} \frac{\delta G_a(t, x)}{\pi g_a(t, x)} = \frac{1}{\pi} \inf_{y \in \mathbb{R}} \frac{\delta G_a(1, y)}{g_a(1, y)}.$$

Let $f(y) = \delta G_a(1, y)/g_a(1, y)$. In the case where $1 < a < 2$ and $|\delta| < 2 - a$, both $\delta G_a(1, y)$ and $g_a(1, y)$ have tails at $\pm\infty$ with polynomial decay of the same rate as $|y|^{-1-a}$: see [27, p. 143] (we use here the fact that $|\delta| \neq 2 - a$). Since $y \mapsto \delta G_a(t, y)$ is unimodal (see [18, Lemma 5.10.1]), we conclude that $f(y) > 0$ for all $y \in \mathbb{R}$, and that $\lim_{y \rightarrow \pm\infty} f(y) > 0$. Therefore, $\inf_{y \in \mathbb{R}} f(y) > 0$, and this completes the proof of Lemma 5.1. □

Lemma 5.2 *Let $f_{b,\nu}(x) = f(x) := (b^2 + x^2)^{-\nu-1/2}$ with $b > 0$ and $\nu \geq 1/2$. Then*

$$\mathcal{F}[f](z) = \int_{\mathbb{R}} dx e^{-izx} f(x) \geq C_\nu b^{-2\nu} \exp(-b|z|), \tag{5.2}$$

for all $b > 0$ and $z \in \mathbb{R}$, where the constant $C_\nu > 0$ is given in (3.18).

Proof Note that the function $f(x)$ is an even function, so its Fourier transform is real-valued, instead of complex-valued, which allows us to bound this transform from below. Indeed, by [14, (7), p. 11], we have that

$$\mathcal{F}[f](z) = \left(\frac{|z|}{b}\right)^\nu \frac{\sqrt{\pi}}{2^\nu \Gamma(\nu + 1/2)} K_\nu(b|z|), \quad \text{for } \Re(b) > 0 \text{ and } \nu > -1/2,$$

where $K_\nu(x)$ is the modified Bessel function of the second kind. Equivalently, we need to prove that the function

$$\mathbb{R}_+ \times \mathbb{R} \ni (b, z) \mapsto \left(\frac{|z|}{b}\right)^\nu \frac{\sqrt{\pi}}{2^\nu \Gamma(\nu + 1/2)} K_\nu(b|z|) b^{2\nu} \exp(b|z|)$$

is uniformly bounded away from zero. By choosing $u = b|z|$, we reduce this problem to bounding the following function

$$\mathbb{R}_+ \ni u \mapsto \frac{\sqrt{\pi}}{2^\nu \Gamma(\nu + 1/2)} h(u) \tag{5.3}$$

away from zero, where $h(u) := u^\nu e^u K_\nu(u)$. By the differential formula for $x^{\pm\nu} K_\nu(x)$ (see, for instance, [20, 51:10:4, p. 532]),

$$h'(u) = e^u u^\nu (K_\nu(u) - K_{\nu-1}(u)).$$

By the integral representation of $K_\nu(z)$ in [21, 10.32.9, p. 252],

$$K_\nu(u) - K_{\nu-1}(u) = \frac{1}{2} \int_0^\infty e^{-u \cosh(t)} \left(e^{\nu t} - e^{-(\nu-1)t} \right) (1 - e^{-t}) dt \geq 0$$

since $\nu \geq 1/2$. Hence, $h'(u) > 0$ and

$$\inf_{u \in \mathbb{R}_+} h(u) = \lim_{u \rightarrow 0} h(u) = 2^{\nu-1} \Gamma(\nu),$$

where we have used the property $K_\nu(u) \sim \frac{1}{2} \Gamma(\nu) (\frac{1}{2}u)^{-\nu}$ as $u \downarrow 0$ (see [21, 10.30.2, p. 252]).

Therefore,

$$C_\nu = \inf_{u \in \mathbb{R}_+} \frac{\sqrt{\pi}}{2^\nu \Gamma(\nu + 1/2)} h(u) = \frac{\Gamma(\nu) \Gamma(1/2)}{2 \Gamma(\nu + 1/2)}.$$

This completes the proof of Lemma 5.2. □

In the next lemma, we gather some properties of the function $g_a(t, x)$.

- Lemma 5.3** (1) For $a > 0$ and $t > 0$, $g_a(t, x - y) \geq \pi 2^{-(a+1)} t^{1/a} g_a(t, x) g_a(t, y)$.
 (2) For $t > 0$ and $z \in \mathbb{R}$, $\mathcal{F}[g_a^2(t, \cdot)](z) \geq C_{a+1/2} \pi^{-2} t^{-1/a} \exp(-t^{1/a}|z|)$.

(3) For all $a > 0$, $t \geq s > 0$ and $x \in \mathbb{R}$, we have

$$\left(g_a^2(t - s, \cdot) * g_a^2(s, \cdot) \right) (x) \geq \frac{C_{a+1/2}^2}{\pi^3 2^{2a+3}} (ts)^{-1/a} (t - s)^{1/a} g_a^2(t - s, x),$$

where the constant $C_{a+1/2}$ is defined in (3.18).

(4) For $t \geq r \geq t/2 > 0$, $g_a(r, x) \geq (t/r)^{1/a} 2^{-1-1/a} g_a(t, x)$.

Proof (1) Because $1 + (u - v)^2 \leq 1 + 2u^2 + 2v^2 \leq (1 + 2u^2)(1 + 2v^2)$, $g_a(t, x - y)$ is bounded from below by $\pi t^{1/a} g_a(t, \sqrt{2} x) g_a(t, \sqrt{2} y)$. Then use the inequality $g_a(t, \sqrt{2} x) \geq 2^{-(a+1)/2} g_a(t, x)$.

(2) We apply Lemma 5.2 with $v = a + 1/2$ and $b = t^{1/a}$.

(3) By (1),

$$\begin{aligned} \left(g_a^2(t - s, \cdot) * g_a^2(s, \cdot) \right) (x) &\geq \pi^2 2^{-2(a+1)} (t - s)^{2/a} g_a^2(t - s, x) \\ &\quad \times \int_{\mathbb{R}} dy g_a^2(t - s, y) g_a^2(s, y). \end{aligned}$$

The integral can be bounded using Plancherel's identity and (2):

$$\begin{aligned} &\int_{\mathbb{R}} dy g_a^2(t - s, y) g_a^2(s, y) \\ &\geq \frac{1}{2\pi} \int_{\mathbb{R}} dz \frac{C_{a+1/2}^2}{\pi^4} ((t - s)s)^{-1/a} \exp\{-|z|((t - s)^{1/a} + s^{1/a})\} \\ &= \frac{C_{a+1/2}^2}{2\pi^5} ((t - s)s)^{-1/a} \frac{2}{(t - s)^{1/a} + s^{1/a}} \\ &\geq \frac{C_{a+1/2}^2}{2\pi^5} ((t - s)s)^{-1/a} t^{-1/a}. \end{aligned}$$

This proves (3).

(4) Notice that for $t \geq r \geq t/2 > 0$,

$$\begin{aligned} g_a(r, x) &= \frac{r^{-1/a}}{\pi} \left(1 + \frac{x^2}{r^{2/a}} \right)^{-(a+1)/2} \geq \frac{r^{-1/a}}{\pi} \left(1 + \frac{x^2}{(t/2)^{2/a}} \right)^{-(a+1)/2} \\ &\geq \frac{r^{-1/a}}{\pi} (t/2)^{1+1/a} ((t/2)^{2/a} + x^2)^{-(a+1)/2} \\ &= \frac{r^{-1/a}}{\pi 2^{1+1/a}} t^{1+1/a} ((t/2)^{2/a} + x^2)^{-(a+1)/2} \geq \frac{r^{-1/a}}{2^{1+1/a}} t^{1/a} g_a(t, x). \end{aligned}$$

□

Proof of Proposition 3.3 Denote

$$(g_a^2)^{\star n}(t, x) := \underbrace{(g_a^2 \star \cdots \star g_a^2)}_{n \text{ factors } g_a^2}(t, x).$$

Notice that by (2.9) and (5.1),

$$\mathcal{K}(t, x; \lambda) = \sum_{n=0}^{\infty} (\lambda^2 {}_{\delta}G_a^2)^{\star(n+1)}(t, x) \geq \sum_{n=0}^{\infty} (\lambda^2 \tilde{C}_{a,\delta}^2 \pi^2 g_a^2)^{\star(n+1)}(t, x). \tag{5.4}$$

We now bound space-time convolutions of g_a^2 with itself. We claim that

$$(\lambda^2 g_a^2)^{\star(n+1)}(t, x) \geq \frac{\lambda^{2(n+1)} \Theta_a^n \Gamma(1/a^*)^{n+1}}{\Gamma((n+1)/a^*)} t^{n/a^*} g_a^2(t, x) \quad \text{for all } n \geq 0, \tag{5.5}$$

where

$$\Theta_a := C_{a+1/2}^2 \pi^{-3} 2^{-2(a+3+1/a)}.$$

The case where $n = 0$ is clear. Consider $n \geq 1$ and assume by induction that (5.5) holds for $n - 1$. By the induction hypothesis and Lemma 5.3(3),

$$\begin{aligned} & (\lambda^2 g_a^2)^{\star(n+1)}(t, x) \\ & \geq \frac{\lambda^{2(n+1)} \Theta_a^{n-1} \Gamma(1/a^*)^n}{\Gamma(n/a^*)} \int_0^t ds (t-s)^{(n-1)/a^*} (g_a^2(t-s, \cdot) * g_a^2(s, \cdot))(x) \\ & \geq K_n t^{-1/a} \int_0^t ds g_a^2(t-s, x) [(t-s)s]^{-1/a} (t-s)^{\frac{n-1}{a^*} + \frac{2}{a}}, \end{aligned} \tag{5.6}$$

where

$$K_n = \frac{\lambda^{2(n+1)} \Theta_a^{n-1} \Gamma(1/a^*)^n C_{a+1/2}^2}{\Gamma(n/a^*) \pi^3 2^{2a+3}}.$$

Notice that $t - s \geq t/2$ for $0 \leq s \leq t/2$, so we apply Lemma 5.3(4) to see that

$$(\lambda^2 g_a^2)^{\star(n+1)}(t, x) \geq \frac{K_n}{2^{2+2/a}} t^{1/a} g_a^2(t, x) \int_0^{t/2} ds [s(t-s)]^{-1/a} (t-s)^{\frac{n-1}{a^*}}. \tag{5.7}$$

For $0 \leq s \leq t/2$, we have $t - s \geq s$, so we replace the last factor $t - s$ by s to see that

$$(\lambda^2 g_a^2)^{\star(n+1)}(t, x) \geq \frac{K_n}{2^{2+2/a}} t^{1/a} g_a^2(t, x) \int_0^{t/2} ds [s(t-s)]^{-1/a} s^{\frac{n-1}{a^*}}.$$

Use the change of variables $s \mapsto t - s$ in (5.7) and add to this last integral to see that

$$\left(\lambda^2 g_a^2\right)^{\star(n+1)}(t, x) \geq \frac{K_n}{2^{2+2/a}} t^{1/a} g_a^2(t, x) \frac{1}{2} \int_0^t ds [s(t-s)]^{-1/a} s^{\frac{n-1}{a^*}}.$$

Then apply Euler’s Beta integral (4.6). This proves (5.5).

Therefore, by (5.4), (5.5) and (3.9),

$$\begin{aligned} \mathcal{K}(t, x; \lambda) &\geq \sum_{n=0}^{+\infty} \left(\lambda^2 \tilde{C}_{a,\delta}^2 \pi^2 g_a^2\right)^{\star(n+1)}(t, x) \\ &\geq \pi^2 \tilde{C}_{a,\delta}^2 \lambda^2 \Gamma(1/a^*) g_a^2(t, x) \sum_{n=0}^{+\infty} \frac{(\lambda^2 \tilde{C}_{a,\delta}^2 \pi^2 \Theta_a \Gamma(1/a^*) t^{1/a^*})^n}{\Gamma((n+1)/a^*)} \\ &= \pi^2 \tilde{C}_{a,\delta}^2 \lambda^2 \Gamma(1/a^*) g_a^2(t, x) E_{1/a^*, 1/a^*}(\lambda^2 \tilde{C}_{a,\delta}^2 \pi^2 \Theta_a \Gamma(1/a^*) t^{1/a^*}). \end{aligned}$$

This proves the statement (3.19) in Proposition 3.3.

As for (3.20), notice that by (3.19),

$$\begin{aligned} (1 \star \mathcal{K})(t, x) &= \int_0^t ds \int_{\mathbb{R}} dy \mathcal{K}(s, y) \\ &\geq C \int_0^t ds E_{1/a^*, 1/a^*} \left(\Upsilon(\lambda, a, \delta) s^{1/a^*}\right) \int_{\mathbb{R}} dy g_a^2(s, y), \end{aligned}$$

where $C = \pi^2 \tilde{C}_{a,\delta}^2 \lambda^2 \Gamma(1/a^*)$. By [21, 5.12.3, p.142] and Euler’s beta integral,

$$\begin{aligned} \int_{\mathbb{R}} dy g_a^2(s, y) &= \frac{s^{-1/a}}{\pi^2} 2 \int_0^\infty dz \frac{1}{(1+z^2)^{a+1}} = \frac{s^{-1/a}}{\pi^2} \int_0^\infty dy \frac{1}{y^{1/2}(1+y)^{a+1}} \\ &= \frac{\Gamma(a+1/2)}{\Gamma(1+a)\pi^{3/2}} s^{-1/a}. \end{aligned}$$

By [22, (1.99), p. 24],

$$\int_0^t ds s^{-1/a} E_{1/a^*, 1/a^*} \left(\Upsilon(\lambda, a, \delta) s^{1/a^*}\right) = t^{1/a^*} E_{1/a^*, 1+1/a^*} \left(\Upsilon(\lambda, a, \delta) t^{1/a^*}\right).$$

This establishes (3.20) and completes the proof of Proposition 3.3. □

5.2 Proofs of Theorems 3.6 and 3.4(2)

Lemma 5.4 *Suppose that $a \in]1, 2[$, $|\delta| < 2 - a$ and $\mu \in \mathcal{M}_{a,+}(\mathbb{R})$, $\mu \neq 0$. Then for all $\epsilon > 0$, there exists a constant C such that for all $t \geq 0$ and $x \in \mathbb{R}$,*

$$(\delta G_a(t, \cdot) * \mu)(x) \geq C 1_{[\epsilon, \infty[}(t) g_a(t, x).$$

Proof Denote $J_0(t, x) = (\delta G_a(t, \cdot) * \mu)(x)$. By the lower bound on $\delta G_a(t, x)$ in (5.1), Lemma 5.3(1) and the scaling property of $g_a(t, x)$, we have

$$\begin{aligned} J_0(t, x) &\geq \tilde{C}_{a,\delta} \pi \int_{\mathbb{R}} \mu(dy) g_a(t, x - y) \\ &\geq \frac{\tilde{C}_{a,\delta} \pi^2}{2^{a+1}} t^{1/a} g_a(t, x) \int_{\mathbb{R}} \mu(dy) g_a(t, y) \\ &= \frac{\tilde{C}_{a,\delta} \pi}{2^{a+1}} g_a(t, x) \int_{\mathbb{R}} \mu(dy) \left(1 + \frac{y^2}{t^{2/a}}\right)^{-\frac{a+1}{2}}. \end{aligned}$$

The above integrand is non-decreasing with respect to t . Hence

$$\begin{aligned} J_0(t, x) &\geq \frac{\tilde{C}_{a,\delta} \pi}{2^{a+1}} 1_{\{t \geq \epsilon\}} g_a(t, x) \int_{\mathbb{R}} \mu(dy) \left(1 + \frac{y^2}{\epsilon^{2/a}}\right)^{-\frac{a+1}{2}} \\ &= \frac{\tilde{C}_{a,\delta} \pi^2 \epsilon^{1/a}}{2^{a+1}} 1_{\{t \geq \epsilon\}} g_a(t, x) \int_{\mathbb{R}} \mu(dy) g_a(\epsilon, y). \end{aligned}$$

Since the function $y \mapsto g_a(\epsilon, y)$ is strictly positive and μ is nonnegative and non-vanishing, the integral is positive. Finally, we can take $C := \tilde{C}_{a,\delta} \pi^2 \epsilon^{1/a} 2^{-(a+1)} \int_{\mathbb{R}} \mu(dy) g_a(\epsilon, y)$. □

Lemma 5.5 *Suppose $\beta > 1$. For all $x \in \mathbb{R}$,*

$$\min_{y \in \mathbb{R}} (|x - y|^\beta + |y|) \geq \begin{cases} \beta^{\frac{\beta}{1-\beta}} + \left| |x| - \beta^{\frac{1}{1-\beta}} \right| & \text{if } |x| \geq \beta^{\frac{1}{1-\beta}}, \\ |x|^\beta & \text{otherwise.} \end{cases}$$

Proof Fix $x \in \mathbb{R}$ and set $f(y) = |x - y|^\beta + |y|$. Assume first that $x \geq 0$. By studying the sign of the derivative of $f'(y)$, we find that if $x \geq \beta^{\frac{1}{1-\beta}}$, then f achieves its minimum at $y = x - \beta^{\frac{1}{1-\beta}}$. If $0 \leq x \leq \beta^{\frac{1}{1-\beta}}$, then f achieves its minimum at 0. The case $x < 0$ is treated similarly. □

Proof of Theorem 3.6 (1) In the following, we use C to denote some nonnegative constant, which may depend on a, δ and L_ρ , and can change from line to line. Fix $p \geq 2$. By (4.21) and (4.22), when $t \geq 1$,

$$(J_0^2 \star \widehat{\mathcal{K}}_p)(t, x) \leq Ct^2 \left(1 + \exp(\widehat{\gamma}_p^{a^*} t)\right) |J_0(t, x)|,$$

where the constant $\widehat{\gamma}_p$ is defined in (3.11). By (3.1) with $\bar{\zeta} = 0$ and (3.22), for $\alpha \geq 0$,

$$\begin{aligned} \lim_{t \rightarrow +\infty} \frac{1}{t} \sup_{|x| \geq \exp(\alpha t)} \log \|u(t, x)\|_p^2 &= \lim_{t \rightarrow +\infty} \frac{1}{t} \sup_{|x| \geq \exp(\alpha t)} \log (J_0^2 \star \widehat{\mathcal{K}}_p)(t, x) \\ &\leq \widehat{\gamma}_p^{a^*} - \alpha\beta. \end{aligned}$$

Now, $\widehat{\gamma}_p^{a^*} - \alpha\beta < 0$ if and only if $\alpha > \beta^{-1} \widehat{\gamma}_p^{a^*}$. Therefore,

$$\bar{e}(p) := \inf \left\{ \alpha > 0 : \lim_{t \rightarrow \infty} \frac{1}{t} \sup_{|x| \geq \exp(\alpha t)} \log \mathbb{E} (|u(t, x)|^p) < 0 \right\} \leq \frac{\widehat{\gamma}_p^{a^*}}{\beta} < +\infty.$$

Concerning the sufficient condition for (3.22), suppose that for some $\eta > 0$, $\int_{\mathbb{R}} |\mu|(dy)(1 + |y|^\eta) < \infty$. We consider first the case where $\eta \in]0, 1 + a[$. Then by (4.3),

$$\begin{aligned} |J_0(t, x)| &\leq \int_{\mathbb{R}} |\mu|(dy) \frac{K_{a,0}(1 + t)}{1 + |x - y|^{1+a}} \\ &\leq CK_{a,0}(1 + t) \sup_{y \in \mathbb{R}} \left[(1 + |y|)(1 + |x - y|)^{(1+a)/\eta} \right]^{-\eta}. \end{aligned}$$

Let $\tilde{\beta} = (1 + a)/\eta > 1$. Notice that

$$(1 + |x - y|^{\tilde{\beta}})(1 + |y|) \geq 1 + |x - y|^{\tilde{\beta}} + |y|.$$

By Lemma 5.5, we see that

$$|J_0(t, x)| \leq \tilde{C}(1 + t) \frac{1}{1 + |x|^\eta},$$

which is condition (3.22) with $\beta = \eta$.

Now consider the case where $\eta \geq 1 + a$. Notice that if $\eta > 1 + a$, then we generally do not expect (3.22) to hold with $\beta = \eta$, since for instance, $J_0(t, x) \sim 1/|x|^{1+a}$ as $|x| \rightarrow \infty$ when $\mu = \delta_0$. Observe that

$$|J_0(t, x)| \leq \int_{\mathbb{R}} \frac{|\mu|(dy)}{1 + |x - y|^{1+a}} \delta G_a(t, x - y) \left(1 + |x - y|^{1+a}\right).$$

From (4.3), we deduce that for $t \geq 1$,

$$\delta G_a(t, x - y) \left(1 + |x - y|^{1+a}\right) \leq K_{a,n} t.$$

Let $\varphi = \eta/(1 + a)$, so that $\varphi \geq 1$. Since for some $\tilde{c} > 0$,

$$\begin{aligned} (1 + |x - y|^2)(1 + |y|^{2\varphi}) &\geq \frac{1}{2} + |x - y|^2 + \frac{1}{2} + |y|^{2\varphi} \geq \tilde{c} \left[1 + |x - y|^2 + |y|^2\right] \\ &\geq \tilde{c} \left(1 + \frac{x^2}{2}\right), \end{aligned}$$

we see that for all $t \geq 1$ and $x \in \mathbb{R}$, there is $c > 0$ such that

$$|J_0(t, x)| \leq CK_{a,n} t \int_{\mathbb{R}} \frac{|\mu|(dy)}{[(1 + |x - y|^2)(1 + |y|^{2\varphi})]^{(1+a)/2}} (1 + |y|^\eta)$$

$$\leq \tilde{C} \frac{t}{(1+x^2)^{(1+a)/2}} \int_{\mathbb{R}} |\mu|(dy) (1+|y|^n),$$

which implies (3.22) with $\beta = 1 + a$.

(2) We only need to consider the case $p = 2$ because $\underline{e}(p) \geq \underline{e}(2)$ for $p \geq 2$. Assume first that $\underline{c} = 0$. Fix any $\epsilon \in]0, t/2[$, choose a constant \widehat{C} according to Lemma 5.4 such that

$$J_0(t, x) = (\delta G_a(t, \cdot) * \mu) \geq \widehat{C} 1_{] \epsilon, \infty[}(t) g_a(t, x) =: I_\epsilon(t, x).$$

By (3.3),

$$\|u(t, x)\|_2^2 \geq J_0^2(t, x) + (J_0^2 \star \underline{K})(t, x) \geq (I_\epsilon^2 \star \underline{K})(t, x).$$

Set $\Upsilon = \Upsilon(l_\rho, a, \delta)$ (see (3.17)). By Proposition 3.3 and Lemma 5.3(3),

$$\begin{aligned} (I_\epsilon^2 \star \underline{K})(t, x) &\geq \widehat{C}^2 C \int_0^{t-\epsilon} ds E_{1/a^*, 1/a^*}(\Upsilon s^{1/a^*}) \int_{\mathbb{R}} dy g_a^2(t-s, x-y) g_a^2(s, y) \\ &\geq \frac{\widehat{C}^2 C C_{a+1/2}^2}{\pi^3 2^{2a+3}} t^{-1/a} \int_0^{t-\epsilon} ds E_{1/a^*, 1/a^*}(\Upsilon s^{1/a^*}) \\ &\quad \times s^{-1/a} (t-s)^{1/a} g_a^2(t-s, x). \end{aligned}$$

Notice that

$$g_a^2(t-s, x) \geq (t-s)^2 t^{-2} g_a^2(t, x).$$

Hence,

$$\begin{aligned} (I_\epsilon^2 \star \underline{K})(t, x) &\geq \frac{\widehat{C}^2 C C_{a+1/2}^2}{\pi^3 2^{2a+3}} t^{-2-1/a} g_a^2(t, x) \int_0^{t-\epsilon} ds \\ &\quad \times E_{1/a^*, 1/a^*}(\Upsilon s^{1/a^*}) s^{-1/a} (t-s)^{2+1/a}. \end{aligned}$$

The above integral is bounded from below by

$$\begin{aligned} E_{1/a^*, 1/a^*}(\Upsilon (t-2\epsilon)^{1/a^*}) \int_{t-2\epsilon}^{t-\epsilon} ds s^{-1/a} (t-s)^{2+1/a} \\ \geq E_{1/a^*, 1/a^*}(\Upsilon (t-2\epsilon)^{1/a^*}) \frac{\epsilon^{2+1/a}}{(t-\epsilon)^{1/a}} \epsilon. \end{aligned}$$

Therefore, we have

$$(I_\epsilon^2 \star \underline{K})(t, x) \geq \bar{C} g_a^2(t, x) t^{-2-1/a} (t-\epsilon)^{-1/a} E_{1/a^*, 1/a^*}(\Upsilon (t-2\epsilon)^{1/a^*}), \tag{5.8}$$

where

$$\bar{C} = \frac{\epsilon^{3+1/a} \widehat{C}^2 C C_{a+1/2}^2}{\pi^3 2^{2a+3}}.$$

Because $x \mapsto g_a(t, x)$ is an even function, decreasing for $x \geq 0$, we deduce that for all $\beta > 0$,

$$\begin{aligned} \sup_{|x| > \exp(\beta t)} \|u(t, x)\|_2^2 &\geq \bar{C} g_a^2(t, \exp(\beta t)) t^{-2-1/a} (t - \epsilon)^{-1/a} \\ &\quad \times E_{1/a^*, 1/a^*} \left(\Upsilon(t - 2\epsilon)^{1/a^*} \right). \end{aligned}$$

Because $a > 0$, there exists $t_0 \geq 0$ such that for all $t \geq t_0$, $t^{2/a} \leq e^{2\beta t}$, so

$$g_a^2(t, \exp(\beta t)) = \frac{1}{\pi^2} \frac{t^2}{(t^{2/a} + e^{2\beta t})^{a+1}} \geq \frac{1}{\pi^2} \frac{t^2}{2^{a+1} e^{2\beta(a+1)t}}.$$

Finally, by the asymptotic expansion of the Mittag-Leffler function in Lemma 4.2,

$$\lim_{t \rightarrow \infty} \frac{1}{t} \sup_{|x| > \exp(\beta t)} \log \|u(t, x)\|_2^2 \geq \Upsilon^{a^*} - 2\beta(a + 1). \tag{5.9}$$

Therefore,

$$\begin{aligned} \underline{e}(2) &= \sup \left\{ \beta > 0 : \lim_{t \rightarrow +\infty} \frac{1}{t} \sup_{|x| > \exp(\beta t)} \log \|u(t, x)\|_2^2 > 0 \right\} \\ &\geq \sup \left\{ \beta > 0 : \Upsilon^{a^*} - 2\beta(a + 1) > 0 \right\} = \frac{\Upsilon^{a^*}}{2(a + 1)}. \end{aligned}$$

Now let us consider the case where there is $c > 0$ with $J_0 \geq c$, or $\underline{c} \neq 0$. In this case, by (3.3) and Proposition 3.3,

$$\|u(t, x)\|_2^2 \geq (c^2 + \underline{c}^2) (1 \star \underline{K})(t, x) \geq C t^{1/a^*} E_{1/a^*, 1+1/a^*} \left(\Upsilon t^{1/a^*} \right).$$

This lower bound does not depend on x and hence, by Lemma 4.2, we get (5.9) with the right-hand side replaced by Υ^{a^*} .

This completes the proof of Theorem 3.6. □

Proof of Theorem 3.4 (2) If $\underline{c} \neq 0$, then from (3.3) and Proposition 3.3, for some constant $C > 0$,

$$\|u(t, x)\|_2^2 \geq \underline{c}^2 (1 \star \underline{K})(t, x) \geq C \underline{c}^2 t^{1/a^*} E_{1/a^*, 1+1/a^*} \left(\Upsilon(l_\rho, a, \delta) t^{1/a^*} \right),$$

where the constant $\Upsilon(l_\rho, a, \delta)$ is defined in (3.17). Then use the asymptotic expansion of $E_{\alpha,\beta}(z)$ in Lemma 4.2 to obtain

$$\underline{m}_2(x) \geq \Upsilon(l_\rho, a, \delta)^{a^*}. \tag{5.10}$$

If $\underline{\zeta} = 0$, then from (3.3), (5.8) and the asymptotics of $E_{\alpha,\beta}(z)$ in Lemma 4.2, we obtain, via the calculation that led to (5.9), but without replacing x by $\exp(\beta t)$, the same lower bound as (5.10). Note that this lower bound does not depend on x . This proves the statement (2) with $p = 2$. For $p > 2$, we use Hölder's inequality

$$\mathbb{E} \left[|u(t, x)|^2 \right] \leq \mathbb{E} \left[|u(t, x)|^p \right]^{2/p}.$$

Hence, $\underline{m}_p(x) \geq \frac{p}{2} \underline{m}_2(x)$. This completes the proof of Theorem 3.4. □

Acknowledgments The authors thank Davar Khoshnevisan for interesting discussions on this problem. A preliminary version of this work was presented at a conference at Michigan State University, East Lansing, USA, August 2013. The authors also thank two anonymous referees for a careful reading of this paper and many useful suggestions, which led to significant improvements.

References

1. Balan, R., Conus, D.: Intermittency for the wave and heat equations with fractional noise in time. *Ann. Probab.* (to appear)
2. Balan, R., Conus, D.: A note on intermittency for the fractional heat equation. *Stat. Probab. Lett.* **95**, 6–14 (2014)
3. Bertini, L., Cancrini, N.: The stochastic heat equation: Feynman-Kac formula and intermittence. *J. Stat. Phys.* **78**(5–6), 1377–1401 (1995)
4. Carlen, E., Krée, P.: L^p estimates on iterated stochastic integrals. *Ann. Probab.* **19**(1), 354–368 (1991)
5. Carmona, R.A., Molchanov, S.A.: Parabolic Anderson problem and intermittency. *Mem. Am. Math. Soc.* **108**(518), Viii+125 (1994)
6. Chen, L., Dalang, R.C.: Moments and growth indices for nonlinear stochastic heat equation with rough initial conditions. *Ann. Probab.* (2015). [arXiv:1307.0600](https://arxiv.org/abs/1307.0600)
7. Chen, L., Dalang, R.C.: Moment bounds and asymptotics for the stochastic wave equation. *Stoch. Process. Appl.* **125**, 1605–1628 (2015)
8. Chen, L., Kim, K.: A comparison principle for the nonlinear stochastic fractional heat equation (2014). [arXiv:1410.0604](https://arxiv.org/abs/1410.0604)
9. Conus, D., Joseph, M., Khoshnevisan, D., Shiu, S.-Y.: Initial measures for the stochastic heat equation. *Ann. Inst. Henri Poincaré Probab. Stat.* **50**(1), 136–153 (2014)
10. Conus, D., Khoshnevisan, D.: On the existence and position of the farthest peaks of a family of stochastic heat and wave equations. *Probab. Theory Relat. Fields* **152**(3–4), 681–701 (2012)
11. Dalang, R.C.: Extending the martingale measure stochastic integral with applications to spatially homogeneous s.p.d.e.'s. *Electron. J. Probab.* **4**(6), 1–29 (1999)
12. Debbi, L.: Explicit solutions of some fractional partial differential equations via stable subordinators. *J. Appl. Math. Stoch. Anal.* **2006**, 93502 (2006). doi:[10.1155/JAMSA/2006/93502](https://doi.org/10.1155/JAMSA/2006/93502)
13. Debbi, L., Dozzi, M.: On the solutions of nonlinear stochastic fractional partial differential equations in one spatial dimension. *Stoch. Process. Appl.* **115**(11), 1764–1781 (2005)
14. Erdélyi, A., Magnus, W., Oberhettinger, F., Tricomi, F.G.: *Tables of Integral Transforms*, vol. I. McGraw-Hill Book Company Inc., New York (1954)
15. Foondun, M., Khoshnevisan, D.: Intermittence and nonlinear parabolic stochastic partial differential equations. *Electron. J. Probab.* **14**(21), 548–568 (2009)
16. Gawronski, W.: On the bell-shape of stable densities. *Ann. Probab.* **12**(1), 230–242 (1984)

17. Khoshnevisan, D.: Analysis of Stochastic Partial Differential Equations. CBMS Regional Conference Series in Mathematics, vol. 119, American Mathematical Society, Providence (2014)
18. Lukacs, E.: Characteristic Functions, 2nd edn. Hafner Publishing Co., New York (1970)
19. Mainardi, F., Luchko, Y., Pagnini, G.: The fundamental solution of the space-time fractional diffusion equation. *Fract. Calc. Appl. Anal.* **4**(2), 153–192 (2001)
20. Oldham, K., Myland, J., Spanier, J.: An Atlas of Functions, 2nd edn. Springer, New York (2009)
21. Olver, F.W.J., Lozier, D.W., Boisvert, R.F., Clark, C.W. (eds.): NIST Handbook of Mathematical Functions. U.S. Department of Commerce National Institute of Standards and Technology, Washington, DC (2010)
22. Podlubny, I.: Fractional Differential Equations. Academic Press Inc., San Diego, CA (1999)
23. Sato, K.: Lévy Processes and Infinitely Divisible Distributions. Revised edition of the 1999 English translation. Cambridge University Press, Cambridge (2013)
24. Uchaikin, V.V., Zolotarev, V.M.: Chance and Stability. VSP, Utrecht (1999)
25. Walsh, J.B.: An Introduction to Stochastic Partial Differential Equations. École d'été de probabilités de Saint-Flour. XIV–1984, pp. 265–439. Springer, Berlin (1986)
26. Yosida, K.: Functional Analysis. Springer, Berlin (1995)
27. Zolotarev, V.M.: One-Dimensional Stable Distributions. American Mathematical Society, Providence (1986)