

Financial Mathematics

MATH 5870/6870¹
Fall 2021

Le Chen

lzc0090@auburn.edu

Last updated on
September 2, 2021

Auburn University
Auburn AL

¹Based on Robert L. McDonald's *Derivatives Markets*, 3rd Ed, Pearson, 2013.

Chapter 20. Brownian Motion and Ito Lemma

Chapter 20. Brownian Motion and Ito Lemma

§ 20.1 The Black-Scholes assumption about stock prices

§ 20.2 Brownian motion

§ 20.3 Geometric Brownian motion

§ 20.4 The Ito formula

§ 20.5 The Sharpe ratio

§ 20.6 Risk-neutral valuation

§ 20.7 Problems

Chapter 20. Brownian Motion and Ito Lemma

§ 20.1 The Black-Scholes assumption about stock prices

§ **20.2 Brownian motion**

§ 20.3 Geometric Brownian motion

§ 20.4 The Ito formula

§ 20.5 The Sharpe ratio

§ 20.6 Risk-neutral valuation

§ 20.7 Problems

Definition 20.2-1 A real-valued stochastic process $Z(t)$ is called a **Brownian motion** or **Wiener process** if

1. It starts at 0:

$$Z(0) = 0.$$

2. For $0 \leq s < t$, the increment $Z(t) - Z(s)$ is normally distributed with mean zero and variance $t - s$:

$$Z(t) - Z(s) \sim N(0, t - s).$$

3. Its increments are independent: if

$$0 \leq t_0 \leq t_1 \leq \dots \leq t_k,$$

then

$$\mathbb{P}(Z(t_i) - Z(t_{i-1}) \in H_i, 1 \leq i \leq k) = \prod_{i=1}^k \mathbb{P}(Z(t_i) - Z(t_{i-1}) \in H_i).$$

Remark 20.2-1 One can always construct a **continuous version** of the Brownian motion; from now on, we always assume that Brownian motion is a continuous process.

Theorem 20.2-1 (Some properties of Brownian motion)

1. $Z(t)$ is **nowhere differentiable**.

(Hence, $dZ(t)$ requires some special treatment.)

2. $Z(t)$ satisfies the **scaling property**:

$$\tilde{Z}(t) := \frac{1}{\sqrt{c}} Z(ct) \text{ is also a B.M. for all } c > 0.$$

3. $Z(t)$ is a **martingale**, namely,

$$\mathbb{E}(Z(t+s)|Z(t)) = Z(t).$$

4. For any $t > 0$, $Z(t) \sim N(0, t)$ and

$$\mathbb{E}(Z(t)Z(s)) = \min(t, s) \quad \text{for all } t, s \geq 0.$$

5. $Z(t)$ is **translation invariant**, namely,

$$\tilde{Z}(t) := Z(t+t_0) - Z(t_0) \text{ is also a B.M. for all } t_0 \geq 0.$$

Proof. Part (1) goes beyond this course. All the rest could be proved using our current knowledge.



Arithmetic Brownian motion

Definition 20.2-2 Let $Z(t)$ be a B.M. Then the process $X(t)$ given by

$$dX(t) = \alpha dt + \sigma dZ(t)$$

is called an **arithmetic Brownian motion**. Equivalently, $X(t)$ can be written in the following integral representation:

$$X(t) = X(0) + \int_0^t \alpha ds + \int_0^t \sigma dZ(s).$$

Remark 20.2-2

1. $X(t)$ is normally distributed:

$$X(t) = \sigma t + \sigma Z(t) \sim N(\sigma t, \sigma^2 t).$$

2. $X(t)$ takes both positive and negative values almost surely.
3. αt is a drift term.

The Ornstein-Uhlenbeck process

Definition 20.2-3 Let $Z(t)$ be a B.M. Then the process $X(t)$ given by

$$dX(t) = \lambda (\alpha - X(t)) dt + \sigma dZ(t)$$

is called the **Ornstein-Uhlenbeck process**.

Remark 20.2-3 Equivalently, $X(t)$ can be written in the following integral representation:

$$X(t) = X(0) + \lambda \int_0^t (\alpha - X(s)) ds + \int_0^t \sigma dZ(s),$$

which is an integral equation (unknown X appears on both sides).

Remark 20.2-4 We have introduced **mean reversion** in the drift term.