

## Euler Formulas

The following series representations can easily be obtained by the standard technique that produces the Maclaurin expansions of functions.

$$e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \frac{x^4}{4!} + \frac{x^5}{5!} + \dots \quad (1)$$

$$\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots \quad (2)$$

$$\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots \quad (3)$$

Observe that if  $i^2 = -1$  then:

$$\begin{aligned} i^1 &= i \\ i^2 &= -1 \\ i^3 &= -i \\ i^4 &= 1 \\ i^5 &= i \\ &\vdots \end{aligned}$$

and the pattern continues. Substituting  $ix$  for  $x$  in equation (1) and also  $-ix$  for  $x$  yields.

$$e^{ix} = 1 + ix - \frac{x^2}{2!} - \frac{ix^3}{3!} + \frac{x^4}{4!} + \frac{ix^5}{5!} - \frac{x^6}{6!} - \frac{ix^7}{7!} + \dots \quad (4)$$

$$e^{-ix} = 1 - ix - \frac{x^2}{2!} + \frac{ix^3}{3!} + \frac{x^4}{4!} - \frac{ix^5}{5!} - \frac{x^6}{6!} + \frac{ix^7}{7!} + \dots \quad (5)$$

Recognizing formulas 2 and 3 as parts of formula 4 and 5 we have:

$$e^{ix} = \cos x + i \sin x \quad (6)$$

$$e^{-ix} = \cos x - i \sin x \quad (7)$$

Solving (6) and (7) for sine and cosine yields:

$$\begin{aligned} \cos x &= \frac{1}{2}(e^{ix} + e^{-ix}) \\ \sin x &= \frac{1}{2i}(e^{ix} - e^{-ix}). \end{aligned}$$

Substituting  $x = \pi$  in equation (6) we get the famous,

$$e^{i\pi} = -1.$$

Recall the definition of the hyperbolic functions and one can see a formal relationship to the trig functions:

$$\begin{aligned}\cosh x &= \frac{1}{2}(e^x + e^{-x}) \\ \sinh x &= \frac{1}{2}(e^x - e^{-x}).\end{aligned}$$

And derive the following

$$\begin{aligned}\cosh x &= \cos(ix) \\ \sinh x &= \frac{1}{i} \sin(ix) = -i \sin(ix).\end{aligned}$$

The hyperbolic trigonometric functions were introduced in the 1760's independently by Vincenzo Riccati and Johann Lambert. They have similar algebraic identities to the usual trig functions; for example:

$$\cosh^2(x) - \sinh^2 x = 1.$$