

Predicate Logic / First Order Logic

The predicate logic system, also called first order logic, subsumes the logical rules that we have already considered (the 'propositional logic'). It adds quantifiers, free variables and relations.

Quantifiers. There are two quantifiers \forall, \exists :

\forall means 'for all'

\exists means 'there exists'.

The letters, x, y, z are typically used for free variables.

Follow are some examples of relations I've made up.

$H(x)$ means x is human

$M(x)$ means x is mortal.

So the statement $H(\text{Socrates})$ is true; and the statement $H(\text{Mickey Mouse})$ is not true. So I can use a quantifier to tell us that there is such a thing as 'Socrates' and that Socrates is human:

$$\exists x(x = \text{Socrates})(H(x)).$$

We would read this as, 'There exists an x such that x is *Socrates* so that x is human. And I can say that humans are mortal:

$$\forall x(H(x) \Rightarrow M(x)).$$

The standard syllogism about Socrates can be written,

$$[(\forall x(H(x) \Rightarrow M(x))) \wedge (\exists x(x = \text{Socrates})(H(x)))] \Rightarrow M(\text{Socrates}).$$

A familiar arithmetic relation is $R(x, y)$ means x is less than y , which may be abbreviated $x < y$. So we can state the tri-part axiom (assume that the variables x and y are numbers; this avoids leading off with the relation stating $\forall x(x \text{ is a number})$):

$$\forall x(\forall y((x < y) \vee (y < x) \vee (x = y))).$$

Exercise 1. Convert each of the following statements into a symbolic representation in the predicate logic. Assume that $R(x, y)$ is some arbitrary relation.

- a.) For some pair x, y the relation $R(x, y)$ is true.
- b.) For some pair x, y the relation $R(x, y)$ is false.
- c.) The relation $R(x, y)$ is always true.
- d.) The relation $R(x, y)$ is always false.
- e.) For some value of x the relation $R(x, y)$ is always true.
- f.) For some value of x the relation $R(x, y)$ is always false.
- g.) For each y there is an x so that $R(x, y)$ true.
- h.) There is an x so that for each y , $R(x, y)$ true.

Exercise 2. Translate the following logical statement into words, assume that all free variables are numbers and that the relations $<, +, -, \times, \dots$ take on the values we're used to. Assume that $0, 1, 3, 3.56$, etc. are the fixed constants with their traditional values. Determine which logical statements are true and find counter examples for those that are not true. Assume that $R(x, y)$ is some arbitrary relation.

- a. $\forall x \forall y (x + y = y + x)$
- b. $\forall x \forall y (x - y = y - x)$
- c. $\forall x \exists y (x + y = y + x)$
- d. $\forall x \exists y (x - y = y - x)$
- e. $\forall x \exists y (x < y)$
- f. $\exists y \forall x (x < y)$
- g. $\forall x ((0 < x) \Rightarrow \exists y ((y < x) \wedge (0 < y)))$
- h. $\sim (\exists y \forall x (x < y))$
- i. $\forall y \exists x (\sim (x < y))$
- j. $\sim (\exists y \forall x R(x, y)) \Leftrightarrow \forall y \exists x (\sim R(x, y))$
- k. $\sim (\exists y \forall x R(x, y)) \Leftrightarrow \exists y \forall x (\sim R(x, y))$.

Examples

For these examples we assume our universe is the set of real numbers. So for a number x , $H(x)$ is a statement about x . For example let $H(x)$ be the

statement $x^2 > 4$. I'll use the notation $H(x) : x^2 > 4$ to indicate this. In this case $H(1.5)$ is false and the statement $H(3)$ is true. The statement $\exists x(H(x))$ is true because there exists an x such that $H(x)$, namely $x = 3$. And the statement $\forall x(H(x))$ is false because $H(x)$ is not true for all x . Therefore I can write, for this meaning of $H(x)$:

$$\begin{aligned} \exists x(H(x)) & \text{ equivalently } \exists x(x^2 > 4) \text{ is true;} \\ \forall x(H(x)) & \text{ equivalently } \forall x(x^2 > 4) \text{ is false.} \end{aligned}$$

Exercise 3. For an arbitrary relation $D(x)$, justify the following :

$$\begin{aligned} \sim (\exists x D(x)) &= \forall x (\sim D(x)); \\ \sim (\forall x D(x)) &= \exists x (\sim D(x)). \end{aligned}$$

When there are two variables it gets more complicated. Suppose $B(x, y)$ is the statement $x < y^2$. The statement

$$\forall x \exists y B(x, y)$$

should be parsed as follows:

$$\forall x (\exists y (B(x, y))).$$

So in order for this statement to be true, we need to have that for all possible x values, the statement $\exists y(B(x, y))$ must be true. We can interpret this in English as

“For all x there is a y so that $B(x, y)$.”

Equivalently:

“If x is a number, there is a y so that $B(x, y)$.”

or

“If x is a number, there is a y so that $x < y^2$.”

This statement is true, in fact any y such that $y > \sqrt{|x|}$ should work. If this is correct, this says that the statement

$$\forall x (B(x, \sqrt{|x|} + 1))$$

is true.

However the statement

$$\exists y \forall x B(x, y)$$

is not true because there is no (fixed) number y so that $x < y^2$ is always true; in fact we can always pick $x = y^2 + 1$ so make the statement false.

Here are how the other possibilities go:

$$\exists x \forall y (x < y^2) \text{ is true;}$$

$$\forall y \exists x (x < y^2) \text{ is true.}$$

Problem: Is the following true for an arbitrary relation $R(x, y)$:

$$\exists x \forall y (R(x, y)) \Rightarrow \forall y \exists x (R(x, y)).$$